SANTA CRUZ BIOTECHNOLOGY, INC.

VHY (G-7): sc-166173



BACKGROUND

Mitogen-activated protein (MAP) kinases are a large class of proteins involved in signal transduction pathways that are activated by a range of stimuli and mediate a number of physiological and pathological changes in the cell. Dual specificity phosphatases (DSPs) are a subclass of the protein tyrosine phosphatase (PTP) gene superfamily, which are selective for dephosphorylating critical phosphothreonine and phosphotyrosine residues within MAP kinases. DSP gene expression is induced by a host of growth factors and/or cellular stresses, thereby negatively regulating MAP kinase superfamily members including MAPK/ERK, SAPK/JNK and p38. VH1-related member Y (VHY) is a member of a subgroup of myristoylated VH1-like small dual specificity phosphatases. It is highly expressed in testis, specifically in pachytene spermatocytes (midstage of meiotic division I) and round spermatids. VHY localizes to the plasma membrane in transfected 293T or NIH/3T3 cells.

REFERENCES

- Keyse, S.M. 1995 An emerging family of dual specificity MAP kinase phosphatases. Biochim. Biophys. Acta 1265: 152-160.
- Sun, H. 1998. Functional studies of dual-specificity phosphatases. Methods Mol. Biol. 84: 307-318.
- Camps, M., et al. 2000. Dual specificity phosphatases: a gene family for control of MAP kinase function. FASEB J. 14: 6-16.
- Alonso, A., et al. 2004. VHY, a novel myristoylated testis-restricted dual specificity protein phosphatase related to VHX. J. Biol. Chem. 279: 32586-32591.

CHROMOSOMAL LOCATION

Genetic locus: DUSP15 (human) mapping to 20q11.21; Dusp15 (mouse) mapping to 2 H1.

SOURCE

VHY (G-7) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 32-61 near the N-terminus of VHY of human origin.

PRODUCT

Each vial contains 200 $\mu g\, lgG_1$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

VHY (G-7) is available conjugated to agarose (sc-166173 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-166173 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-166173 PE), fluorescein (sc-166173 FITC), Alexa Fluor[®] 488 (sc-166173 AF488), Alexa Fluor[®] 546 (sc-166173 AF546), Alexa Fluor[®] 594 (sc-166173 AF594) or Alexa Fluor[®] 647 (sc-166173 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-166173 AF680) or Alexa Fluor[®] 790 (sc-166173 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-166173 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

VHY (G-7) is recommended for detection of VHY of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

VHY (G-7) is also recommended for detection of VHY in additional species, including canine, bovine, porcine and avian.

Suitable for use as control antibody for VHY siRNA (h): sc-61786, VHY siRNA (m): sc-61787, VHY shRNA Plasmid (h): sc-61786-SH, VHY shRNA Plasmid (m): sc-61787-SH, VHY shRNA (h) Lentiviral Particles: sc-61786-V and VHY shRNA (m) Lentiviral Particles: sc-61787-V.

Molecular Weight of VHY: 26 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210, F9 cell lysate: sc-2245 or VHY (h): 293T Lysate: sc-116612.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



VHY (G-7): sc-166173. Western blot analysis of VHY expression in non-transfected: sc-117752 (**A**) and human VHY transfected: sc-116612 (**B**) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.