

AP-4 (H-5): sc-166216

BACKGROUND

AP-2 transcription factor family members include AP-2 α , AP-2 β and AP-2 γ , which specifically bind to the DNA consensus sequence CCCAGGC and initiate transcription of selected genes. AP-2, also known as ERF-1, plays a role in regulating estrogen receptor expression. AP-2 β , a splice variant of AP-2 α , inhibits AP-2 activity. Besides subscribing to the AP-2 complex, AP-2 α , AP-2 β and AP-2 γ proteins compose the OB2-1 transcription factor complex. OB2-1 specifically upregulates expression of the proto-oncogene c-ErbB-2, which is overexpressed in 25-30% of breast cancers. The gene encoding AP-2 α maps to human chromosome 6p24.3. AP-2 α may play an important role in the development of ectodermal-derived tissues. Deleterious mutations involving the AP-2 α gene are linked to microphthalmia, corneal clouding and other anterior eye chamber defects. The ubiquitously expressed AP-4 transcription factor specifically binds to the DNA consensus sequence 5'-CAGCTG-3'. AP-4 interacts with promoters for immunoglobulin- κ gene families and simian virus 40. AP-4 may enhance the transcription of the human Huntington's disease gene. AP-4 is a helix-loop-helix protein that contains two distinctive leucine repeat elements.

REFERENCES

- Williams, T., et al. 1988. Cloning and expression of AP-2, a cell-type-specific transcription factor that activates inducible enhancer elements. *Genes Dev.* 2: 1557-1569.
- Hu, Y.F., et al. 1990. Transcription factor AP-4 contains multiple dimerization domains that regulate dimer specificity. *Genes Dev.* 4: 1741-1752.
- Moser, M., et al. 1995. Cloning and characterization of a second AP-2 transcription factor: AP-2 β . *Development* 121: 2779-2788.
- Bosher, J.M., et al. 1996. A family of AP-2 proteins regulates c-ErbB-2 expression in mammary carcinoma. *Oncogene* 13: 1701-1707.

CHROMOSOMAL LOCATION

Genetic locus: TFAP4 (human) mapping to 16p13.3; Tcfap4 (mouse) mapping to 16 A1.

SOURCE

AP-4 (H-5) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 2-43 at the N-terminus of AP-4 of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-166216 X, 200 μ g/0.1 ml.

Blocking peptide available for competition studies, sc-166216 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

AP-4 (H-5) is recommended for detection of AP-4 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

AP-4 (H-5) is also recommended for detection of AP-4 in additional species, including canine.

Suitable for use as control antibody for AP-4 siRNA (h): sc-37690, AP-4 siRNA (m): sc-37691, AP-4 shRNA Plasmid (h): sc-37690-SH, AP-4 shRNA Plasmid (m): sc-37691-SH, AP-4 shRNA (h) Lentiviral Particles: sc-37690-V and AP-4 shRNA (m) Lentiviral Particles: sc-37691-V.

AP-4 (H-5) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of AP-4: 48 kDa.

Positive Controls: MOLT-4 nuclear extract: sc-2151, AP-4 (m): 293T Lysate: sc-118454 or TK-1 whole cell lysate: sc-364798.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



AP-4 (H-5): sc-166216. Western blot analysis of AP-4 expression in MOLT-4 (A) and WEHI-231 (B) nuclear extracts and TK-1 whole cell lysate (C).



AP-4 (H-5): sc-166216. Western blot analysis of AP-4 expression in non-transfected: sc-117752 (A) and mouse AP-4 transfected: sc-118454 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

- Kang, H.G., et al. 2021. Genetic variants in histone modification regions are associated with the prognosis of lung adenocarcinoma. *Sci. Rep.* 11: 21520.

RESEARCH USE

For research use only, not for use in diagnostic procedures.