

DDX41 (E-8): sc-166255

BACKGROUND

DDX41 (probable ATP-dependent RNA helicase DDX41, DEAD box protein abstract homolog) is a 622 amino acid protein encoded by the human gene DDX41. DDX41 belongs to the DEAD box helicase family (DDX41 subfamily) and contains one CCHC-type zinc finger, one helicase ATP-binding domain and one helicase C-terminal domain. DDX41 is required during post-transcriptional gene expression and is thought to be involved in pre-mRNA splicing. DDX41 is believed to be a probable ATP-dependent RNA helicase. RNA helicases are highly conserved enzymes that utilize the energy derived from NTP hydrolysis to modulate the structure of RNA. RNA helicases participate in all biological processes that involve RNA, including transcription, splicing and translation.

REFERENCES

1. Irion, U. and Leptin, M. 2000. Developmental and cell biological functions of the *Drosophila* DEAD-box protein abstract. *Curr. Biol.* 9: 1373-1381.
2. Abdul-Ghani, M., Hartman, K.L. and Ngsee, J.K. 2005. Abstract interacts with and regulates the expression of sorting nexin-2. *J. Cell. Physiol.* 204: 210-218.
3. Sekito, A., Taira, T., Niki, T., Iguchi-Arigo, S.M. and Ariga, H. 2005. Stimulation of transforming activity of DJ-1 by abstract, a DJ-1-binding protein. *Int. J. Oncol.* 26: 685-689.
4. Abdelhaleem, M. 2005. RNA helicases: regulators of differentiation. *Clin. Biochem.* 38: 499-503.
5. Nousiainen, M., Sillje, H.H., Sauer, G., Nigg, E.A. and Körner, R. 2006. Phosphoproteome analysis of the human mitotic spindle. *Proc. Natl. Acad. Sci. USA* 103: 5391-5396.
6. Savitsky, M., Kwon, D., Georgiev, P., Kalmykova, A. and Gvozdev, V. 2006. Telomere elongation is under the control of the RNAi-based mechanism in the *Drosophila* germline. *Genes Dev.* 20: 345-354.

CHROMOSOMAL LOCATION

Genetic locus: DDX41 (human) mapping to 5q35.3; Ddx41 (mouse) mapping to 13 B1.

SOURCE

DDX41 (E-8) is a mouse monoclonal antibody raised against amino acids 111-206 mapping within an internal region of DDX41 of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-166255 X, 200 µg/0.1 ml.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

DDX41 (E-8) is recommended for detection of DDX41 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

DDX41 (E-8) is also recommended for detection of DDX41 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for DDX41 siRNA (h): sc-91765, DDX41 siRNA (m): sc-142938, DDX41 shRNA Plasmid (h): sc-91765-SH, DDX41 shRNA Plasmid (m): sc-142938-SH, DDX41 shRNA (h) Lentiviral Particles: sc-91765-V and DDX41 shRNA (m) Lentiviral Particles: sc-142938-V.

DDX41 (E-8) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of DDX41: 70 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210, Jurkat whole cell lysate: sc-2204 or DDX41 (h): 293 Lysate: sc-113244.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



DDX41 (E-8): sc-166255. Western blot analysis of DDX41 expression in non-transfected: sc-110760 (A) and human DDX41 transfected: sc-113244 (B) 293T whole cell lysates.



DDX41 (E-8): sc-166255. Western blot analysis of DDX41 expression in NIH/3T3 whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.