

HAX-1 (N-17): sc-16636

BACKGROUND

HAX-1 (HS1-associated protein X-1 or HS1-binding protein protein X-1), encodes a novel protein. HAX-1 has previously been shown to associate with HS1, a protein specifically expressed in cells of the hematopoietic lineage, and is thought to be involved in signal transduction in B cells and apoptosis. Though first identified as a protein that associates with HS1, recent data has also revealed interactions between HAX-1 and three disparate proteins: Polycystin-2 (derived from the PKD2 gene), a protein linked to polycystic kidney disease; Cortactin; and EBNA-LP (Epstein-Barr virus nuclear antigen leader protein). Additionally, HAX-1 has been identified as a binding partner to the carboxy-terminus of the K15 protein of Kaposi's sarcoma-associated herpesvirus. K15 interacts with cellular HAX-1 *in vitro* and *in vivo*. Furthermore, HAX-1 co-localizes with K15 in the endoplasmic reticulum and mitochondria. Immunofluorescence experiments show that in most cells PKD2 and HAX-1 co-localize in the cell body, but in some cells PKD2 and HAX-1 also are sorted into cellular processes and lamellipodia. The HAX-1 gene is expressed ubiquitously among tissues. Its protein is localized mainly in mitochondria, but also in endoplasmic reticulum and the nuclear envelope of the cell.

CHROMOSOMAL LOCATION

Genetic locus: HAX1 (human) mapping to 1q21.3; Hax1 (mouse) mapping to 3 F1.

SOURCE

HAX-1 (N-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of HAX-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-16636 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

HAX-1 (N-17) is recommended for detection of HAX-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

HAX-1 (N-17) is also recommended for detection of HAX-1 in additional species, including porcine.

Suitable for use as control antibody for HAX-1 siRNA (h): sc-43365, HAX-1 siRNA (m): sc-43366, HAX-1 shRNA Plasmid (h): sc-43365-SH, HAX-1 shRNA Plasmid (m): sc-43366-SH, HAX-1 shRNA (h) Lentiviral Particles: sc-43365-V and HAX-1 shRNA (m) Lentiviral Particles: sc-43366-V.


Molecular Weight of HAX-1: 35 kDa.

Positive Controls: A-673 cell lysate: sc-2414, Saos-2 cell lysate: sc-2235 or MES-SA/Dx5 cell lysate: sc-2284.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



HAX-1 (N-17): sc-16636. Immunoperoxidase staining of formalin fixed, paraffin-embedded human rectum tissue showing cytoplasmic staining of glandular cells.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **HAX-1 (B-11): sc-166845**, our highly recommended monoclonal alternative to HAX-1 (N-17).