

# Troponin T-FS (C-12): sc-166533

## BACKGROUND

Actin is a highly conserved protein that is expressed in all eukaryotic cells. Actin filaments can form both stable and labile structures and are crucial components of microvilli and the contractile apparatus of muscle cells. Myosin is a hexamer of two heavy chains (MHC) and four light chains (MLC) that interacts with actin to generate the force for diverse cellular movements, including cytokinesis, phagocytosis and muscle contraction. Troponin facilitates the interaction between actin and myosin by binding to calcium. Troponin is made up of at least two subunits, which are divergent in cardiac muscle, fast skeletal muscle and slow skeletal muscle. Structures of skeletal muscle Troponin are composed of Troponin C (the sensor), Troponin I (the regulator) and Troponin T (the link to the muscle thin filament). Troponin C is dumbbell-shaped and has a hydrophobic pocket that increases the contractile force of muscle fibers. Troponin C has two isoforms: fast and slow. Fast Troponin C has two calcium binding sites while slow/cardiac Troponin C has a single calcium binding site.

## REFERENCES

1. Parmacek, M.S. and Leiden, J.M. 1989. Structure and expression of the murine slow/cardiac Troponin C gene. *J. Biol. Chem.* 264: 13217-13225.
2. Koppe, R.I., Hallauer, P.L., Karpati, G. and Hastings, K.E. 1989. cDNA clone and expression analysis of rodent fast and slow skeletal muscle Troponin I mRNAs. *J. Biol. Chem.* 264: 14327-14333.
3. Ausoni, S., Campione, M., Picard, A., Moretti, P., Vitadello, M., DeNardi, C. and Schiaffino, S. 1994. Structure and regulation of the mouse cardiac Troponin I gene. *J. Biol. Chem.* 269: 339-346.
4. Potter, J.D., Sheng, Z., Pan, B.S. and Zhao, J. 1995. A direct regulatory role for troponin T and a dual role for Troponin C in the Ca<sup>2+</sup> regulation of muscle contraction. *J. Biol. Chem.* 270: 2557-2562.
5. Barkalow, K. and Hartwig, J.H. 1995. Actin cytoskeleton. Setting the pace of cell movement. *Curr. Biol.* 5: 1000-1002.
6. Baker, J.P. and Titus, M.A. 1998. Myosins: matching functions with motors. *Curr. Opin. Cell Biol.* 10: 80-86.

## CHROMOSOMAL LOCATION

Genetic locus: TNNT3 (human) mapping to 11p15.5; Tnnt3 (mouse) mapping to 7 F5.

## SOURCE

Troponin T-FS (C-12) is a mouse monoclonal antibody raised against amino acids 36-258 mapping at the C-terminus of fast skeletal muscle Troponin T of human origin.

## PRODUCT

Each vial contains 200 µg IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

Troponin T-FS (C-12) is recommended for detection of fast skeletal muscle Troponin T of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Troponin T-FS siRNA (h): sc-36742, Troponin T-FS siRNA (m): sc-36743, Troponin T-FS shRNA Plasmid (h): sc-36742-SH, Troponin T-FS shRNA Plasmid (m): sc-36743-SH, Troponin T-FS shRNA (h) Lentiviral Particles: sc-36742-V and Troponin T-FS shRNA (m) Lentiviral Particles: sc-36743-V.


Molecular Weight of Troponin T-FS: 31 kDa.

Positive Controls: human skeletal muscle extract: sc-363776, mouse skeletal muscle extract: sc-364250 or rat skeletal muscle extract: sc-364810.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



Troponin T-FS (C-12): sc-166533. Western blot analysis of Troponin T-FS expression in human skeletal muscle (A) and mouse skeletal muscle (B) tissue extracts.



Troponin T-FS (C-12): sc-166533. Western blot analysis of Troponin T-FS expression in rat skeletal muscle tissue extract.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.