

▶ ILT (D-6): sc-166549

BACKGROUND

Leukocyte immunoglobulin-like receptors (ILTs, also known as LIRs) are members of the immunoglobulin superfamily of glycoproteins and are predominantly expressed by monocytes, B cells, dendritic cells, natural killer (NK) cells, peripheral blood leukocytes and tissues such as placenta, lung and liver. There are several members of the ILT family, including ILT-1, ILT-2, ILT-3, ILT-4, ILT-5, ILT-6, ILT-7, ILT-8, ILT-11, LIR-6 and LIR-8. These ILT proteins are divided into two subfamilies, namely subfamily A (ILT-1, ILT-6, ILT-7, ILT-8, ILT-11 and LIR-6) and subfamily B (ILT-2, ILT-3, ILT-4, ILT-5 and LIR-8), the former of which function as stimulating receptors and the latter of which function as inhibitory receptors. Characteristically, members of subfamily A have transmembrane regions containing a charged arginine residue through which they initiate stimulatory cascades, while members of subfamily B contain cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs) through which they induce inhibitory signaling cascades.

REFERENCES

1. Samaridis, J. and Colonna, M. 1997. Cloning of novel immunoglobulin superfamily receptors expressed on human myeloid and lymphoid cells: structural evidence for new stimulatory and inhibitory pathways. *Eur. J. Immunol.* 27: 660-665.
2. Cosman, D., et al. 1997. A novel immunoglobulin superfamily receptor for cellular and viral MHC class I molecules. *Immunity* 7: 273-282.
3. Colonna, M., et al. 1997. A common inhibitory receptor for major histocompatibility complex class I molecules on human lymphoid and myelomonocytic cells. *J. Exp. Med.* 186: 1809-1818.
4. André, P., et al. 2001. New nomenclature for MHC receptors. *Nat. Immunol.* 2: 661.
5. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604811. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Foster, C.E., et al. 2003. Crystal structure of the human natural killer (NK) cell activating receptor NKp46 reveals structural relationship to other leukocyte receptor complex immunoreceptors. *J. Biol. Chem.* 278: 46081-46086.
7. Shiroishi, M., et al. 2003. Human inhibitory receptors Ig-like transcript 2 (ILT2) and ILT4 compete with CD8 for MHC class I binding and bind preferentially to HLA-G. *Proc. Natl. Acad. Sci. USA* 100: 8856-8861.

CHROMOSOMAL LOCATION

Genetic locus: LILRA2/LILRB1/LILRB2/LILRB3/LILRA3 (human) mapping to 19q13.42.

SOURCE

ILT (D-6) is a mouse monoclonal antibody raised against amino acids 162-461 mapping within an extracellular domain of ILT-2 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

ILT (D-6) is recommended for detection of ILT family members 1, 2, 4, 5 and 6 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Molecular Weight of different ILT proteins: 33-71 kDa.

Positive Controls: ILT-4 (h): 293T Lysate: sc-115693 or human spleen extract: sc-363779.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



ILT (D-6): sc-166549. Western blot analysis of ILT-4 expression in non-transfected: sc-117752 (A) and human ILT-4 transfected: sc-115693 (B) 293T whole cell lysates.



ILT (D-6): sc-166549. Western blot analysis of ILT expression in human spleen tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.