SANTA CRUZ BIOTECHNOLOGY, INC.

group VI iPLA₂ (E-8): sc-166616



BACKGROUND

Phospholipases catalyze the release of fatty acids from phospholipids. One member of the phospholipase family, iPLA₂, is detected as a membrane-bound protein with multiple smaller isoforms, which result from alternative splicing. Two isoforms, Ankyrin-iPLA₂-1 and -2, lack the catalytic domain and are thought to be involved in the negative regulation of iPLA₂ activity. The SH-iPLA₂ isoform is cytoplasmically localized. The human gene encoding iPLA₂ maps to chromosome 22q13.1. Another phopholipase, iPLA₂, belongs to a family of secretory phospholipases A₂, which represent an expanding family of related enzymes. iPLA₂ has both membrane bound and secreted forms that are encoded by a single gene. iPLA₂ is involved in the regulation of phospholipid metabolism in biomembranes and in eicosanoid biosynthesis.

REFERENCES

- 1. Scott, D.L., et al. 1991. Structures of free and inhibited human secretory phospholipase A₂ from inflammatory exudate. Science 254: 1007-1010.
- 2. Lehninger, A., et al. 1993. Principles of Biochemistry Second Edition. New York: Worth Publishers.

CHROMOSOMAL LOCATION

Genetic locus: PLA2G6 (human) mapping to 22q13.1; Pla2g6 (mouse) mapping to 15 E1.

SOURCE

group VI iPLA₂ (E-8) is a mouse monoclonal antibody raised against amino acids 1-120 of group VI iPLA₂ of human origin.

PRODUCT

Each vial contains 200 μg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

group VI iPLA₂ (E-8) is recommended for detection of calcium-independent PLA₂ of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for group VI iPLA₂ siRNA (h): sc-43819, group VI iPLA₂ siRNA (m): sc-43820, group VI iPLA₂ siRNA (r): sc-270117, group VI iPLA₂ shRNA Plasmid (h): sc-43819-SH, group VI iPLA₂ shRNA Plasmid (m): sc-43820-SH, group VI iPLA₂ shRNA Plasmid (r): sc-270117-SH, group VI iPLA₂ shRNA (h) Lentiviral Particles: sc-43819-V, group VI iPLA₂ shRNA (m) Lentiviral Particles: sc-43820-V and group VI iPLA₂ shRNA (r) Lentiviral Particles: sc-270117-V.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG K BP-HRP: sc-516102 or m-IgG K BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-lgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA





group VI iPLA₂ (E-8): sc-166616. Western blot analysis of group VI iPLA₂ expression in F9 (**A**), PC-12 (**B**) and c4 (C) whole cell lysates and mouse liver tiss extract (D)

group VI iPLA2 (E-8): sc-166616. Immunofluorescence staining of methanol-fixed NIH/3T3 cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human small intestine tissue showing cytoplasmic staining of glandular cells (B).

SELECT PRODUCT CITATIONS

- 1. Jiao, L., et al. 2015. Deficiency of group VIA phospholipase A_2 (iPLA₂ β) renders susceptibility for chemical-induced colitis. Dig. Dis. Sci. 60: 3590-3602.
- 2. Nelson, A.J., et al. 2020. Macrophage polarization is linked to Ca²⁺⁻ independent phospholipase $A_2\beta$ -derived lipids and cross-cell signaling in mice. J. Lipid Res. 61: 143-158.
- 3. Jin, T., et al. 2021. iPLA₂ β contributes to ER stress-induced apoptosis during myocardial ischemia/reperfusion injury. Cells 10: 1446.
- 4. Zhao, Z., et al. 2021. Lipid metabolism is a novel and practical source of potential targets for antiviral discovery against porcine parvovirus. Vet. Microbiol. 261: 109177.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Molecular Weight of group VI iPLA₂: 88 kDa.