

ASB-9 (E-4): sc-166737

BACKGROUND

Members of the suppressor of cytokine signaling (SOCS) family of proteins contain C-terminal regions of homology called the SOCS box, which serves to couple SOCS proteins and their binding partners with the Elongin B and C complex. Several other families of proteins also contain SOCS boxes, but differ from the SOCS proteins in the type of domain they contain upstream of the SOCS box. The largest family of SOCS box-containing proteins is the Ankyrin repeat and SOCS box-containing (ASB) protein family. Members of the ASB family include ASB-1 through ASB-18 and are involved in a variety of biological processes. ASB-9 is a 294 amino acid member of this family. It contains six Ankyrin repeats and one SOCS box domain. ASB-9 functions as a ubiquitin ligase and, via its SOCS box domain, it specifically interacts with creatine kinase-B, targeting it for degradation and regulating its expression within the cell. Two isoforms exist for ASB-9 due to alternative splicing events.

REFERENCES

1. Bork, P. 1993. Hundreds of Ankyrin-like repeats in functionally diverse proteins: mobile modules that cross phyla horizontally? *Proteins* 17: 363-374.
2. Hilton, D.J., et al. 1998. Twenty proteins containing a C-terminal SOCS box form five structural classes. *Proc. Natl. Acad. Sci. USA* 95: 114-119.
3. Kile, B.T., et al. 2000. Cloning and characterization of the genes encoding the Ankyrin repeat and SOCS box-containing proteins ASB-1, ASB-2, ASB-3 and ASB-4. *Gene* 258: 331-341.
4. Kile, B.T., et al. 2002. The SOCS box: a tale of destruction and degradation. *Trends Biochem. Sci.* 27: 235-241.
5. Kohroki, J., et al. 2005. ASB proteins interact with Cullin5 and Rbx2 to form E3 ubiquitin ligase complexes. *FEBS Lett.* 579: 6796-6802.

CHROMOSOMAL LOCATION

Genetic locus: Asb9 (mouse) mapping to X F5.

SOURCE

ASB-9 (E-4) is a mouse monoclonal antibody raised against amino acids 1-70 mapping at the N-terminus of ASB-9 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

ASB-9 (E-4) is available conjugated to agarose (sc-166737 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-166737 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-166737 PE), fluorescein (sc-166737 FITC), Alexa Fluor® 488 (sc-166737 AF488), Alexa Fluor® 546 (sc-166737 AF546), Alexa Fluor® 594 (sc-166737 AF594) or Alexa Fluor® 647 (sc-166737 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-166737 AF680) or Alexa Fluor® 790 (sc-166737 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

ASB-9 (E-4) is recommended for detection of ASB-9 of mouse and rat origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ASB-9 siRNA (m): sc-141293, ASB-9 shRNA Plasmid (m): sc-141293-SH and ASB-9 shRNA (m) Lentiviral Particles: sc-141293-V.

Molecular Weight of ASB-9: 32 kDa.


Positive Controls: mouse testis extract: sc-2405 or Hep G2 cell lysate: sc-2227.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



ASB-9 (E-4): sc-166737. Near-infrared western blot analysis of ASB-9 expression in mouse testis tissue extract. Blocked with UltraCruz® Blocking Reagent: sc-516214. Detection reagent used: m-IgGκ BP-CFL 680: sc-516180.



ASB-9 (E-4): sc-166737. Western blot analysis of ASB-9 expression in mouse testis tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.