

NCoA-3 (E-11): sc-166751

BACKGROUND

Nuclear receptors for steroids, thyroid hormones and retinoic acids are ligand-dependent transcription factors that activate transcription through specific DNA binding sites in their target genes. Several related transcriptional co-activators and corepressors have been described that work in concert with the steroid receptor family to either induce or repress transcription from hormone-responsive elements. This family includes GRIP1 (for GR interacting protein 1), also designated NCoA-2 or TIF-2; SRC-1 (for steroid receptor co-activator-1), also designated NCoA-1; and NCoA-3, also designated RAC3, ACTR, AIB-1 (for amplified in breast cancer) and p/CIP (for p300/CBP/co-integrator protein), which displays elevated expression in estrogen receptor positive ovarian and breast cancers and is required for the transcriptional activation of p300/CBP-dependent transcription factors.

REFERENCES

1. Ribeiro, R.C., et al. 1995. The nuclear hormone receptor gene superfamily. *Annu. Rev. Med.* 46: 443-453.
2. Onate, S.A., et al. 1995. Sequence and characterization of a co-activator for the steroid hormone receptor superfamily. *Science* 270: 1354-1357.
3. Hong, H., et al. 1996. GRIP1, a novel mouse protein that serves as a transcriptional co-activator in yeast for the hormone binding domains of steroid receptors. *Proc. Natl. Acad. Sci. USA* 93: 4948-4952.
4. Li, H., et al. 1997. RAC3, a steroid/nuclear receptor-associated co-activator that is related to SRC-1 and TIF2. *Proc. Natl. Acad. Sci. USA* 94: 8479-8484.
5. Anzick, S.L., et al. 1997. AIB1, a steroid receptor co-activator amplified in breast and ovarian cancer. *Science* 277: 965-968.

CHROMOSOMAL LOCATION

Genetic locus: NCOA3 (human) mapping to 20q13.12; Ncoa3 (mouse) mapping to 2 H3.

SOURCE

NCoA-3 (E-11) is a mouse monoclonal antibody raised against amino acids 1321-1400 of AIB-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-166751 X, 200 µg/0.1 ml.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

NCoA-3 (E-11) is recommended for detection of NCoA-3 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NCoA-3 siRNA (h): sc-29636, NCoA-3 siRNA (m): sc-29637, NCoA-3 shRNA Plasmid (h): sc-29636-SH, NCoA-3 shRNA Plasmid (m): sc-29637-SH, NCoA-3 shRNA (h) Lentiviral Particles: sc-29636-V and NCoA-3 shRNA (m) Lentiviral Particles: sc-29637-V.

NCoA-3 (E-11) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of NCoA-3: 160 kDa.

Positive Controls: HeLa nuclear extract: sc-2120, MCF7 whole cell lysate: sc-2206 or K-562 whole cell lysate: sc-2203.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



NCoA-3 (E-11): sc-166751. Western blot analysis of NCoA-3 expression in HeLa nuclear extract (A) and MCF7 (B) and NIH/3T3 (C) whole cell lysates.



NCoA-3 (E-11): sc-166751. Western blot analysis of NCoA-3 expression in Jurkat (A) and K-562 (B) whole cell lysates.

SELECT PRODUCT CITATIONS

1. Wang, D., et al. 2019. C/EBPδ-Slug-Lox1 axis promotes metastasis of lung adenocarcinoma via oxLDL uptake. *Oncogene*. E-published.

RESEARCH USE

For research use only, not for use in diagnostic procedures.