

# eIF2B $\gamma$ (D-8): sc-166768

## BACKGROUND

The initiation of protein synthesis in eukaryotic cells is regulated by interactions between protein initiation factors and RNA molecules. The eukaryotic initiation complex eIF2B exists as a five subunit complex composed of eIF2B $\alpha$ , eIF2B $\beta$ , eIF2B $\gamma$ , eIF2B $\delta$  and eIF2B $\epsilon$ . The eIF2B complex catalyzes the exchange of GDP for GTP on the eIF2 complex, following the interaction of eIF2/GTP with the 40S ribosomal subunit. Guanine nucleotide exchange factor (GEF) activity is exhibited by the eIF2B $\epsilon$  subunit alone, but is greater in the presence of all five eIF2B subunits. Phosphorylation of eIF2 inhibits GEF activity of eIF2B, an inhibition that requires the eIF2B $\alpha$  subunit.

## REFERENCES

- Henderson, R.A., et al. 1994. The  $\delta$ -subunit of murine guanine nucleotide exchange factor eIF2B. Characterization of cDNAs predicts isoforms differing at the amino-terminal end. *J. Biol. Chem.* 269: 30517-30523.
- Flowers, K.M., et al. 1995. Structure and sequence of the gene encoding the  $\alpha$ -subunit of rat translation initiation factor 2B. *Biochim. Biophys. Acta* 1264: 163-167.
- Price, N.T., et al. 1996. Cloning of cDNA for the  $\gamma$ -subunit of mammalian translation initiation factor 2B, the guanine nucleotide-exchange factor for eukaryotic initiation factor 2. *Biochem. J.* 318: 631-636.
- Price, N.T., et al. 1996. eIF2B, the guanine nucleotide-exchange factor for eukaryotic initiation factor 2. Sequence conservation between the  $\alpha$ ,  $\beta$  and  $\delta$  subunits of eIF2B from mammals and yeast. *Biochem. J.* 318: 637-643.
- Asuru, A.I., et al. 1996. Cloning and characterization of cDNAs encoding the  $\epsilon$ -subunit of eukaryotic initiation factor 2B from rabbit and human. *Biochim. Biophys. Acta* 1307: 309-317.
- Webb, B.L., et al. 1997. Eukaryotic initiation factor 2B (eIF2B). *Int. J. Biochem. Cell Biol.* 29: 1127-1131.
- Fabian, J.R., et al. 1997. Subunit assembly and guanine nucleotide exchange activity of eukaryotic initiation factor 2B expressed in Sf9 cells. *J. Biol. Chem.* 272: 12359-12365.

## CHROMOSOMAL LOCATION

Genetic locus: EIF2B3 (human) mapping to 1p34.1.

## SOURCE

eIF2B $\gamma$  (D-8) is a mouse monoclonal antibody raised against amino acids 153-452 mapping at the C-terminus of eIF2B $\gamma$  of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

eIF2B $\gamma$  (D-8) is recommended for detection of eIF2B $\gamma$  of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for eIF2B $\gamma$  siRNA (h): sc-35274, eIF2B $\gamma$  shRNA Plasmid (h): sc-35274-SH and eIF2B $\gamma$  shRNA (h) Lentiviral Particles: sc-35274-V.


Molecular Weight of eIF2B $\gamma$ : 50 kDa.

Positive Controls: K-562 nuclear extract: sc-2130, HeLa nuclear extract: sc-2120 or IMR-32 nuclear extract: sc-2148.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG $\kappa$  BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohisto-mount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



eIF2B $\gamma$  (D-8): sc-166768. Western blot analysis of eIF2B $\gamma$  expression in IMR-32 (A), HeLa (B) and K-562 (C) nuclear extracts and HL-60 (D) and RAW 264.7 (E) whole cell lysates. Detection reagent used: m-IgG $\kappa$  BP-HRP: sc-516102.



eIF2B $\gamma$  (D-8): sc-166768. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human duodenum tissue showing cytoplasmic staining of glandular cells (B).

## SELECT PRODUCT CITATIONS

- Moon, S.L. and Parker, R. 2018. eIF2B2 mutations in vanishing white matter disease hypersuppress translation and delay recovery during the integrated stress response. *RNA* 24: 841-852.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.