

IL-19 (N-15): sc-16727

BACKGROUND

IL-14 (high-molecular-weight B-cell growth factor or HMW-BCGF) is a 53kDa protein that plays a significant role in the rapid proliferation of aggressive B-cell type non-Hodgkin's lymphoma. IL-19 and IL-20 are IL-10-related cytokines. IL-19 is involved in melanoma differentiation, and IL-20 plays an essential role in epidermal function and psoriasis. Like other interleukin molecules, IL-21 is a small secreted molecule, and it has potent effects on lymphoid cells. IL-21 is most closely related to IL-2 and IL-15 and has a role in the proliferation and maturation of natural killer cell populations from bone marrow, in the proliferation of mature B-cell populations co-stimulated with anti-CD40, and in the proliferation of T cells co-stimulated with anti-CD3.

REFERENCES

1. Ford, R., Tamayo, A., Martin, B., Niu, K., Claypool, K., Cabanillas, F. and Ambrus, J. Jr. 1995. Identification of B-cell growth factors (interleukin-14; high molecular weight-B-cell growth factors) in effusion fluids from patients with aggressive B-cell lymphomas. *Blood* 86: 283-293.
2. Gallagher, G., Dickensheets, H., Eskdale, J., Izotova, L.S., Mirochnitchenko, O.V., Peat, J.D., Vazquez, N., Pestka, S., Donnelly, R.P. and Kotenko, S.V. 2000. Cloning, expression and initial characterization of interleukin-19 (IL-19), a novel homologue of human interleukin-10 (IL-10). *Genes Immun.* 1: 442-450.
3. Parrish-Novak, J., Dillon, S.R., Nelson, A., Hammond, A., Sprecher, C., Gross, J.A., Johnston, J., Madden, K., Xu, W., West, J., Schrader, S., Burkhead, S., Heipel, M., Brandt, C., Kuijper, J.L., Kramer, J., Conklin, D., Presnell, S.R., Berry, J., Shiota, F., et al. 2000. Interleukin 21 and its receptor are involved in NK cell expansion and regulation of lymphocyte function. *Nature* 408: 57-63.
4. Blumberg, H., Conklin, D., Xu, W.F., Grossmann, A., Brender, T., Carollo, S., Eagan, M., Foster, D., Haldeman, B.A., Hammond, A., Haugen, H., Jelinek, L., Kelly, J.D., Madden, K., Maurer, M.F., Parrish-Novak, J., Prunkard, D., Sexson, S., Sprecher, C., Waggle, K., et al. 2001. Interleukin 20: discovery, receptor identification, and role in epidermal function. *Cell* 104: 9-19.
5. Vosshenrich, C.A. and Di Santo, J.P. 2001. Cytokines: IL-21 joins the γ (c)-dependent network? *Curr. Biol.* 11: R175-R177.

CHROMOSOMAL LOCATION

Genetic locus: IL19 (human) mapping to 1q32.1.

SOURCE

IL-19 (N-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of IL-19 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-16727 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

IL-19 (N-15) is recommended for detection of IL-19 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

IL-19 (N-15) is also recommended for detection of IL-19 in additional species, including canine.

Suitable for use as control antibody for IL-19 siRNA (h): sc-39659, IL-19 shRNA Plasmid (h): sc-39659-SH and IL-19 shRNA (h) Lentiviral Particles: sc-39659-V.

Molecular Weight of IL-19: 18 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.