

CCDC85B (S-14): sc-167420

BACKGROUND

CCDC85B (coiled-coil domain-containing protein 85B), also known as DIPA (δ -interacting protein A), is a 202 amino acid nuclear and cytoplasmic protein that belongs to the CCDC85 family. As a transcriptional repressor, CCDC85B may inhibit the activity of CTNNB1 in a TP53-dependent manner and thus regulate cell growth. CCDC85B may also play a role in adipocyte differentiation, negatively regulating mitotic clonal expansion. While it is up-regulated by doxorubicin, CCDC85B also interacts with a number of proteins including C/EBP β , C/EBP δ , MSP58, TCF-4 and HDAG (viral phosphoprotein hepatitis δ antigen). The interaction between CCDC85B and HDAG affects viral genomic replication in intact cells. The gene that encodes CCDC85B consists of approximately 1,232 bases and maps to human chromosome 11q13.1.

REFERENCES

1. Brazas, R. and Ganem, D. 1996. A cellular homolog of hepatitis δ antigen: implications for viral replication and evolution. *Science* 274: 90-94.
2. Long, M., de Souza, S.J. and Gilbert, W. 1997. δ -interacting protein A and the origin of hepatitis δ antigen. *Science* 276: 824-825.
3. Online Mendelian Inheritance in Man, OMIM[™]. 2000. Johns Hopkins University, Baltimore, MD. MIM Number: 605360. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Bezy, O., Elabd, C., Cochet, O., Petersen, R.K., Kristiansen, K., Dani, C., Ailhaud, G. and Amri, E.Z. 2005. δ -interacting protein A, a new inhibitory partner of CCAAT/enhancer-binding protein β , implicated in adipocyte differentiation. *J. Biol. Chem.* 280: 11432-11438.
5. Du, X., Wang, Q., Hirohashi, Y. and Greene, M.I. 2006. DIPA, which can localize to the centrosome, associates with p78/MCRS1/MSP58 and acts as a repressor of gene transcription. *Exp. Mol. Pathol.* 81: 184-190.
6. Iwai, A., Hijikata, M., Hishiki, T., Isono, O., Chiba, T. and Shimotohno, K. 2008. Coiled-coil domain containing 85B suppresses the β -catenin activity in a p53-dependent manner. *Oncogene* 27: 1520-1526.
7. Bianchi, D.W., Maron, J.L. and Johnson, K.L. 2010. Insights into fetal and neonatal development through analysis of cell-free RNA in body fluids. *Early Hum. Dev.* 86: 747-752.

CHROMOSOMAL LOCATION

Genetic locus: CCDC85B (human) mapping to 11q13.1; Ccdc85b (mouse) mapping to 19 A.

SOURCE

CCDC85B (S-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CCDC85B of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-167420 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CCDC85B (S-14) is recommended for detection of CCDC85B of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with CCDC85A.

CCDC85B (S-14) is also recommended for detection of CCDC85B in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for CCDC85B siRNA (h): sc-96589, CCDC85B siRNA (m): sc-142150, CCDC85B shRNA Plasmid (h): sc-96589-SH, CCDC85B shRNA Plasmid (m): sc-142150-SH, CCDC85B shRNA (h) Lentiviral Particles: sc-96589-V and CCDC85B shRNA (m) Lentiviral Particles: sc-142150-V.

Molecular Weight of CCDC85B: 22 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.