

CD163L1 (C-14): sc-167427

BACKGROUND

CD163L1 (CD163 molecule-like 1), also known as scavenger receptor cysteine-rich type 1 protein M160 or CD163B, is a 1,453 amino acid protein that belongs to the scavenger receptor cysteine-rich (SRCR) superfamily. Members of the SRCR family contain SRCR domains which are used to influence ligand binding and protein-protein interaction, and are found in cells of the immune system where they exist as membrane-anchored or secreted proteins. Containing twelve SRCR domains, CD163L1 undergoes alternative splicing events to produce three isoforms. CD163L1 isoforms 1 and 2 are single-pass type I membrane proteins, unlike isoform 3, which is a secreted protein. CD163L1 isoform 1 is abundantly expressed in lymph node, spleen, thymus and fetal liver, and isoform 2 is found exclusively in spleen. The gene encoding CD163L1 maps to human chromosome 12p13.31.

REFERENCES

1. Stover, C.M., Schleypen, J., Grønland, J., Speicher, M.R., Schwaeble, W.J. and Holmskov, U. 2000. Assignment of CD163B, the gene encoding M160, a novel scavenger receptor, to human chromosome 12p13.3 by in situ hybridization and somatic cell hybrid analysis. *Cytogenet. Cell Genet.* 90: 246-247.
2. Gronlund, J., Vitved, L., Lausen, M., Skjodt, K. and Holmskov, U. 2000. Cloning of a novel scavenger receptor cysteine-rich type I transmembrane molecule (M160) expressed by human macrophages. *J. Immunol.* 165: 6406-6415.
3. Zhang, Z. and Henzel, W.J. 2004. Signal peptide prediction based on analysis of experimentally verified cleavage sites. *Protein Sci.* 13: 2819-2824.
4. Chen, R., Jiang, X., Sun, D., Han, G., Wang, F., Ye, M., Wang, L. and Zou, H. 2009. Glycoproteomics analysis of human liver tissue by combination of multiple enzyme digestion and hydrazide chemistry. *J. Proteome Res.* 8: 651-661.
5. Online Mendelian Inheritance in Man, OMIM[™]. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 606079: World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Van Gorp, H., Van Breedam, W., Van Doorselaere, J., Delpitte, P.L. and Nauwynck, H.J. 2010. Identification of the CD163 protein domains involved in infection of the porcine reproductive and respiratory syndrome virus. *J. Virol.* 84: 3101-3105.

CHROMOSOMAL LOCATION

Genetic locus: CD163L1 (human) mapping to 12p13.31.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

SOURCE

CD163L1 (C-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a C-terminal cytoplasmic domain of CD163L1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-167427 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CD163L1 (C-14) is recommended for detection of CD163L1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with CD163.

Suitable for use as control antibody for CD163L1 siRNA (h): sc-96130, CD163L1 shRNA Plasmid (h): sc-96130-SH and CD163L1 shRNA (h) Lentiviral Particles: sc-96130-V.

Molecular Weight of CD163L1 isoforms: 159/155/80 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.