

## FATE1 (T-16): sc-167853

### BACKGROUND

FATE1 (fetal and adult testis expressed 1), also known as FATE, CT43 (cancer/testis antigen 43) or BJ-HCC-2, is a cancer/testis antigen found in testis and tumor tissues (specifically hepatocarcinoma cells). Its expression is regulated by SF-1 (steroidogenic factor 1) and WT1 (Wilms' tumor protein), two proteins involved in tumorigenesis, suggesting a role for FATE1 in tumor development. FATE1 is exclusively expressed in testis of the 6-11 week old fetus (around the period of gonadal sex differentiation). At seven weeks, FATE1 is coexpressed with SRY and may play a role in early testicular differentiation. In adults, although predominantly expressed in testis, FATE1 can also be found in heart, kidney, brain, lung and adrenal gland. Mutations in the gene encoding FATE1 may be contributing factors in male infertility.

### REFERENCES

- Olesen, C., et al. 2001. Human FATE is a novel X-linked gene expressed in fetal and adult testis. *Mol. Cell. Endocrinol.* 184: 25-32.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 300450. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Olesen, C., et al. 2003. Mutational analysis of the human FATE gene in 144 infertile men. *Hum. Genet.* 113: 195-201.
- Dong, X.Y., et al. 2003. Identification of two novel CT antigens and their capacity to elicit antibody response in hepatocellular carcinoma patients. *Br. J. Cancer* 89: 291-297.
- Yang, X.A., et al. 2004. Purification and refolding of a novel cancer/testis antigen BJ-HCC-2 expressed in the inclusion bodies of *Escherichia coli*. *Protein Expr. Purif.* 33: 332-338.
- Yang, X.A., et al. 2005. Immunohistochemical analysis of the expression of FATE/BJ-HCC-2 antigen in normal and malignant tissues. *Lab. Invest.* 85: 205-213.
- Doghman, M., et al. 2007. Increased steroidogenic factor-1 dosage triggers adrenocortical cell proliferation and cancer. *Mol. Endocrinol.* 21: 2968-2987.

### CHROMOSOMAL LOCATION

Genetic locus: FATE1 (human) mapping to Xq28.

### SOURCE

FATE1 (T-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of FATE1 of human origin.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-167853 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### APPLICATIONS

FATE1 (T-16) is recommended for detection of FATE1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for FATE1 siRNA (h): sc-90989, FATE1 shRNA Plasmid (h): sc-90989-SH and FATE1 shRNA (h) Lentiviral Particles: sc-90989-V.

Molecular Weight of FATE1: 21 kDa.

Positive Controls: Human testis tissue extract.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.