

GnT-IVA (N-14): sc-168023

BACKGROUND

GnT-IVA (N-acetylglucosaminyltransferase IVA), also known as GlcNAc-T IVA, GnT-IV or MGAT4A (mannosyl (α -1,3-)-glycoprotein β -1,4-N-acetylglucosaminyltransferase, isozyme A) is a type II single-pass membrane protein that belongs to the glycosyltransferase 54 family of proteins. Localizing to the membrane of the Golgi apparatus, GnT-IVA is expressed in thymus, pancreas, prostate, small intestine, lymph node, spleen and peripheral blood leukocytes. It functions as a glycosyltransferase and participates in protein modification by catalyzing the transfer of N-acetylglucosamine (GlcNAc) to mannose residues of N-linked glycans, thereby regulating the formation of tri- and multi-antennary structures. GnT-IVA may be involved in regulating cell differentiation, oncogenesis and the availability of serum glycoproteins and is known to play a role in the development of choriocarcinoma. In addition, GnT-IVA is recognized as a genetic marker for pancreatic cancer as its expression is downregulated in these cancer tissues.

REFERENCES

1. Takamatsu, S., Oguri, S., Minowa, M.T., Yoshida, A., Nakamura, K., Takeuchi, M. and Kobata, A. 1999. Unusually high expression of N-acetylglucosaminyltransferase-IVA in human choriocarcinoma cell lines: a possible enzymatic basis of the formation of abnormal biantennary sugar chain. *Cancer Res.* 59: 3949-3953.
2. Fukuta, K., Abe, R., Yokomatsu, T., Minowa, M.T., Takeuchi, M., Asanagi, M. and Makino, T. 2001. The widespread effect of β 1,4-galactosyltransferase on N-glycan processing. *Arch. Biochem. Biophys.* 392: 79-86.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604623. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Ohtsubo, K., Takamatsu, S., Minowa, M.T., Yoshida, A., Takeuchi, M. and Marth, J.D. 2005. Dietary and genetic control of glucose transporter 2 glycosylation promotes Insulin secretion in suppressing diabetes. *Cell* 123: 1307-1321.
5. Oguri, S., Yoshida, A., Minowa, M.T. and Takeuchi, M. 2006. Kinetic properties and substrate specificities of two recombinant human N-acetylglucosaminyltransferase-IV isozymes. *Glycoconj. J.* 23: 473-480.
6. Thorens, B. 2006. A toggle for type 2 diabetes? *N. Engl. J. Med.* 354: 1636-1638.
7. Ide, Y., Miyoshi, E., Nakagawa, T., Gu, J., Tanemura, M., Nishida, T., Ito, T., Yamamoto, H., Kozutsumi, Y. and Taniguchi, N. 2006. Aberrant expression of N-acetylglucosaminyltransferase-IVa and IVb (GnT-IVa and b) in pancreatic cancer. *Biochem. Biophys. Res. Commun.* 341: 478-482.

CHROMOSOMAL LOCATION

Genetic locus: MGAT4A (human) mapping to 2q11.2; Mgat4a (mouse) mapping to 1 B.

SOURCE

GnT-IVA (N-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of GnT-IVA of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-168023 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

GnT-IVA (N-14) is recommended for detection of GnT-IVA of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with GnT-IVA soluble form; non cross-reactive with GnT-IVB or GnT-IVH.

GnT-IVA (N-14) is also recommended for detection of GnT-IVA in additional species, including equine, bovine, porcine and avian.

Suitable for use as control antibody for GnT-IVA siRNA (h): sc-94905, GnT-IVA siRNA (m): sc-145663, GnT-IVA shRNA Plasmid (h): sc-94905-SH, GnT-IVA shRNA Plasmid (m): sc-145663-SH, GnT-IVA shRNA (h) Lentiviral Particles: sc-94905-V and GnT-IVA shRNA (m) Lentiviral Particles: sc-145663-V.

Molecular Weight of GnT-IVA: 62 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **GnT-IVA (M-71): sc-100785**, our highly recommended monoclonal alternative to GnT-IVA (N-14).