

group IB sPLA₂ (P-13): sc-168045

BACKGROUND

Phospholipase A₂s (PLA₂s) constitute a family of esterases that hydrolyze the sn-2-acyl ester bond in glycerophospholipid molecules. These enzymes are generally calcium-dependent and have been found both intra- and extracellularly. By hydrolyzing the sn-2 bond in glycerophospholipids, PLA₂s release fatty acids. One such fatty acid, arachidonic acid, generates the substrates for the initiation of the arachidonic acid cascade that produces various eicosanoids, many of which are potent mediators of inflammation. As a member of the PLA₂ family, group IB sPLA₂, also known as PLA2G1B (phospholipase A₂, group IB), is a 148 amino acid secreted protein that is expressed in the pancreas. Group IB sPLA₂ catalyzes the reaction of phosphatidylcholine and water into 1-acylglycerophosphocholine and carboxylate, and may play a role in obesity and Insulin resistance.

REFERENCES

- Seilhmer, J.J., et al. 1986. Pancreatic phospholipase A₂: isolation of the human gene and cDNAs from porcine pancreas and human lung. *DNA* 5: 519-527.
- Seilhmer, J.J., et al. 1989. Novel gene exon homologous to pancreatic phospholipase A₂: sequence and chromosomal mapping of both human genes. *J. Cell. Biochem.* 39: 327-337.
- Ishizaki, J., et al. 1999. Cloning and characterization of novel mouse and human secretory phospholipase A₂s. *J. Biol. Chem.* 274: 24973-24979.
- Shakhov, A.N., et al. 2000. SPLASH (PLA2IID), a novel member of phospholipase A₂ family, is associated with lymphotoxin deficiency. *Genes Immun.* 1: 191-199.
- Eckey, R., et al. 2004. Increased hepatic cholesterol accumulation in transgenic mice overexpressing human secretory phospholipase A₂ group IIA. *Inflammation* 28: 59-65.
- Takabatake, N., et al. 2005. A novel polymorphism in secretory phospholipase A₂-IID is associated with body weight loss in chronic obstructive pulmonary disease. *Am. J. Respir. Crit. Care Med.* 172: 1097-1104.
- Lindbom, J., et al. 2005. Interferon γ -induced gene expression of the novel secretory phospholipase A₂ type IID in human monocyte-derived macrophages is inhibited by lipopolysaccharide. *Inflammation* 29: 108-117.
- Online Mendelian Inheritance in Man, OMIM[™]. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 172410. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Igarashi, A., et al. 2009. Gly80Ser polymorphism of phospholipase A₂-IID is associated with cytokine inducibility in A549 cells. *Respiration* 78: 312-321.

CHROMOSOMAL LOCATION

Genetic locus: PLA2G1B (human) mapping to 12q24.31.

SOURCE

group IB sPLA₂ (P-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of group IB sPLA₂ of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-168045 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA). support products.

APPLICATIONS

group IB sPLA₂ (P-13) is recommended for detection of group IB sPLA₂ of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for group IB sPLA₂ siRNA (h): sc-96105, group IB sPLA₂ shRNA Plasmid (h): sc-96105-SH and group IB sPLA₂ shRNA (h) Lentiviral Particles: sc-96105-V.

Molecular Weight of group IB sPLA₂: 16 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and