

KCNF1 (G-15): sc-168237

BACKGROUND

KCNF1 (Potassium voltage-gated channel subfamily F member 1) is a multi-pass membrane-bound protein that acts as an ion channel and is generally expressed as a heterotetramer of potassium channeling proteins. Formerly known as kH1, KCNF1 is usually found as a heteromer with three other potassium channel proteins, KCNG3, KV6.3 and KCNV2. As a potassium channel protein, KCNF1 plays a role in regulating apoptosis and proliferation of pulmonary artery smooth muscle (PASM) cells. Bone morphogenetic proteins (BMPs) restrict proliferation and can induce apoptosis in normal human PASM cells and will upregulate expression of KCNF1 in PASM cells *in vitro*. KCNF1 is expressed in heart, brain, liver, skeletal muscle, kidney and pancreas.

REFERENCES

1. Su, K., Kyaw, H., Fan, P., Zeng, Z., Shell, B.K., Carter, K.C. and Li, Y. 1997. Isolation, characterization, and mapping of two human potassium channels. *Biochem. Biophys. Res. Commun.* 241: 675-681.
2. Lubec, G. and Sohn, S.Y. 2003. RNA microarray analysis of channels and transporters in normal and fetal Down syndrome (Trisomy 21) brain. *J. Neural. Transm. Suppl.* 67: 215-224.
3. Fantozzi, I., Platoshyn, O., Wong, A.H., Zhang, S., Remillard, C.V., Furtado, M.R., Petrauskene, O.V. and Yuan, J.X. 2006. Bone morphogenetic protein-2 upregulates expression and function of voltage-gated K⁺ channels in human pulmonary artery smooth muscle cells. *Am. J. Physiol. Lung Cell Mol. Physiol.* 291: L993-1004.
4. Hassan, M.J., Santos, R.L., Rafiq, M.A., Chahrour, M.H., Pham, T.L., Wajid, M., Hijab, N., Wambangco, M., Lee, K., Ansar, M., Yan, K., Ahmad, W. and Leal, S.M. 2006. A novel autosomal recessive non-syndromic hearing impairment locus (DFNB47) maps to chromosome 2p25.1-p24.3. *Hum. Genet.* 118: 605-610.
5. Judge, S.I., Smith, P.J., Stewart, P.E. and Bever, C.T. 2007. Potassium channel blockers and openers as CNS neurologic therapeutic agents. *Recent Patents CNS Drug Discov.* 2: 200-228.
6. Knight, H.M., Maclean, A., Irfan, M., Naeem, F., Cass, S., Pickard, B.S., Muir, W.J., Blackwood, D.H. and Ayub, M. 2008. Homozygosity mapping in a family presenting with schizophrenia, epilepsy and hearing impairment. *Eur. J. Hum. Genet.* 16: 750-758.

CHROMOSOMAL LOCATION

Genetic locus: KCNF1 (human) mapping to 2p25.1; Kcnf1 (mouse) mapping to 12 A1.1.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

SOURCE

KCNF1 (G-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a C-terminal cytoplasmic domain of KCNF1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-168237 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

KCNF1 (G-15) is recommended for detection of KCNF1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for KCNF1 siRNA (h): sc-94734, KCNF1 siRNA (m): sc-146358, KCNF1 shRNA Plasmid (h): sc-94734-SH, KCNF1 shRNA Plasmid (m): sc-146358-SH, KCNF1 shRNA (h) Lentiviral Particles: sc-94734-V and KCNF1 shRNA (m) Lentiviral Particles: sc-146358-V.

Molecular Weight of KCNF1: 14 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.