

FGF-19 (N-20): sc-16831

BACKGROUND

Fibroblast growth factor-1 (FGF-1), also designated acidic FGF, and fibroblast growth factor-2 (FGF-2), also designated basic FGF, are members of a family of growth factors that stimulate proliferation of cells of mesenchymal, epithelial and neuroectodermal origin. Additional members of the FGF family include the oncogenes FGF-3 (Int2) and FGF-4 (hst/Kaposi), FGF-5, FGF-6, FGF-7 (KGF), FGF-8 (AIGF), FGF-9 (GAF) and FGF-10–FGF-23. Members of the FGF family share 30-55% amino acid sequence identity and similar gene structure, and are capable of transforming cultured cells when overexpressed in transfected cells. Cellular receptors for FGFs are members of a second multigene family including four tyrosine kinases, designated Flg (FGFR-1), Bek (FGFR-L), TKF and FGFR-3.

REFERENCES

- Moore, R., et al. 1986. Sequence, topography and protein coding potential of mouse Int-2: a putative oncogene activated by mouse mammary tumor virus. *EMBO J.* 5: 919-924.
- Delli Bovi, P., et al. 1987. An oncogene isolated by transfection of Kaposi's sarcoma DNA encodes a growth factor that is a member of the FGF family. *Cell* 50: 729-737.
- Zhan, X., et al. 1988. The human FGF-5 oncogene encodes a novel protein related to fibroblast growth factors. *Mol. Cell. Biol.* 8: 3487-3495.
- Rifkin, D.B., et al. 1989. Recent developments in the cell biology of fibroblast growth factor. *J. Cell Biol.* 109: 1-6.
- Marics, I., et al. 1989. Characterization of the HST-related FGF-6 gene, a new member of the fibroblast growth factor gene family. *Oncogene* 4: 335-340.
- Dionne, C.A., et al. 1990. Cloning and expression of two distinct high-affinity receptors cross-reacting with acidic and basic fibroblast growth factors. *EMBO J.* 9: 2685-2692.
- Tanaka, A., et al. 1992. Cloning and characterization of an androgen-induced growth factor essential for the androgen-dependent growth of mouse mammary carcinoma cells. *Proc. Natl. Acad. Sci. USA* 89: 8928-8932.
- Miyamoto, M., et al. 1993. Molecular cloning of a novel cytokine cDNA encoding the ninth member of the fibroblast growth factor family, which has a unique secretion property. *Mol. Cell. Biol.* 13: 4251-4259.

CHROMOSOMAL LOCATION

Genetic locus: FGF19 (human) mapping to 11q13.3.

SOURCE

FGF-19 (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of FGF-19 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-16831 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

FGF-19 (N-20) is recommended for detection of FGF-19 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for FGF-19 siRNA (h): sc-39480, FGF-19 shRNA Plasmid (h): sc-39480-SH and FGF-19 shRNA (h) Lentiviral Particles: sc-39480-V.


Molecular Weight of FGF-19: 24 kDa.

Positive Controls: Y79 cell lysate: sc-2240 or Hep G2 cell lysate: sc-2227.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



FGF-19 (N-20): sc-16831. Western blot analysis of FGF-19 expression in Y79 (A) and Hep G2 (B) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **FGF-19 (H-12): sc-390621** or **FGF-19 (W12): sc-73984**, our highly recommended monoclonal alternatives to FGF-19 (N-20).