SANTA CRUZ BIOTECHNOLOGY, INC.

LOC130951 (K-14): sc-168445



BACKGROUND

The second largest human chromosome, 2 consists of 237 million bases encoding over 1,400 genes and making up approximately 8% of the human genome. A number of genetic diseases are linked to genes on chromosome 2. Harlequin icthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene. The lipid metabolic disorder sitosterolemia is associated with ABCG5 and ABCG8. An extremely rare recessive genetic disorder, Alström syndrome is due to mutations in the ALMS1 gene. Interestingly, chromosome 2 contains what appears to be a vestigial second centromere and vestigial telomeres which gives credence to the hypothesis that human chromosome 2 is the result of an ancient fusion of two ancestral chromosomes seen in modern form today in apes. The LOC130951 gene product has been provisionally designated LOC130951 pending further characterization.

REFERENCES

- Ijdo, J.W., et al. 1991. Origin of human chromosome 2: an ancestral telomere-telomere fusion. Proc. Natl. Acad. Sci. USA 88: 9051-9055.
- 2. Avarello, R., et al. 1992. Evidence for an ancestral alphoid domain on the long arm of human chromosome 2. Hum. Genet. 89: 247-249.
- 3. Hillier, L.W., et al. 2005. Generation and annotation of the DNA sequences of human chromosomes 2 and 4. Nature 434: 724-731.
- 4. Thomas, A.C., et al. 2006. ABCA12 is the major harlequin ichthyosis gene. J. Invest. Dermatol. 126: 2408-2413.
- Akiyama, M., et al. 2007. Compound heterozygous ABCA12 mutations including a novel nonsense mutation underlie harlequin ichthyosis. Dermatology 215: 155-159.
- Marshall, J.D., et al. 2007. Alström syndrome. Eur. J. Hum. Genet. 15: 1193-1202.
- Marshall, J.D., et al. 2007. Spectrum of ALMS1 variants and evaluation of genotype-phenotype correlations in Alström syndrome. Hum. Mutat. 28: 1114-1123.
- 8. Tabas, I. 2007. A two-carbon switch to sterol-induced autophagic death. Autophagy 3: 38-41.
- 9. Wang, D.Q. 2007. Regulation of intestinal cholesterol absorption. Annu. Rev. Physiol. 69: 221-248.

CHROMOSOMAL LOCATION

Genetic locus: C2orf65 (human) mapping to 2p13.1; D6Mm5e (mouse) mapping to 6 C3.

SOURCE

LOC130951 (K-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of LOC130951 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-168445 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

LOC130951 (K-14) is recommended for detection of LOC130951 of human origin, D6Mm5e of mouse origin and corresponding rat homolog by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

LOC130951 (K-14) is also recommended for detection of LOC130951 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for LOC130951 siRNA (h): sc-94504, D6Mm5e siRNA (m): sc-142846, LOC130951 shRNA Plasmid (h): sc-94504-SH, D6Mm5e shRNA Plasmid (m): sc-142846-SH, LOC130951 shRNA (h) Lentiviral Particles: sc-94504-V and D6Mm5e shRNA (m) Lentiviral Particles: sc-142846-V.

Molecular Weight of LOC130951: 59 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluo-rescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.