

striatin (N-20): sc-16856

BACKGROUND

Striatin, SG2NA, and zinedin, the three mammalian members of the striatin family, are multimodular, WD-repeat and calmodulin-binding proteins. Zinedin and SG2NA share with striatin identical protein-protein interaction domains and the same overall domain structure. All three proteins are both cytosolic and membrane-bound and bind calmodulin in the presence of calcium. Striatin is a neuronal, intracellular protein strictly expressed in the somatodendritic compartment, including spines, subsets of neurons, and is considered as a marker of neuronal polarity. Downregulation of striatin, which is expressed in a few subsets of neurons, impairs the growth of dendrites as well as rat locomotor activity. Zinedin is mainly expressed in the central nervous system, whereas SG2NA is mainly expressed in the brain and muscle.

REFERENCES

1. Castets, F., et al. 1996. A novel calmodulin-binding protein, belonging to the WD-repeat family, is localized in dendrites of a subset of CNS neurons. *J. Cell Biol.* 134: 1051-1062.
2. Kachidian, P., et al. 1998. Relationships between striatin-containing neurons and cortical or thalamic afferent fibers in the rat striatum: an ultrastructural study by dual labeling. *Neuroscience* 85: 111-122.
3. Salin, P., et al. 1998. Distribution of striatin, a newly identified calmodulin-binding protein in the rat brain: an *in situ* hybridization and immunocytochemical study. *J. Comp. Neurol.* 397: 41-59.
4. Bartoli, M., et al. 1999. Down-regulation of striatin, a neuronal calmodulin-binding protein, impairs rat locomotor activity. *J. Neurobiol.* 40: 234-243.

CHROMOSOMAL LOCATION

Genetic locus: STRN (human) mapping to 2p22.2; Strn (mouse) mapping to 17 E3.

SOURCE

striatin (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of striatin of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-16856 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

striatin (N-20) is recommended for detection of striatin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for striatin siRNA (h): sc-37649, striatin siRNA (m): sc-37650, striatin shRNA Plasmid (h): sc-37649-SH, striatin shRNA Plasmid (m): sc-37650-SH, striatin shRNA (h) Lentiviral Particles: sc-37649-V and striatin shRNA (m) Lentiviral Particles: sc-37650-V.


Molecular Weight of striatin: 110 kDa.

Positive Controls: A549 cell lysate: sc-2413, SK-N-SH cell lysate: sc-2410 or rat brain extract: sc-2392.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



striatin (N-20): sc-16856. Western blot analysis of striatin expression in A549 (A), C2C12 (B) and SK-N-SH (C) whole cell lysates and rat brain tissue extract (D).



striatin (N-20): sc-16856. Western blot analysis of striatin expression in NIH/3T3 whole cell lysate.

SELECT PRODUCT CITATIONS

1. Strack, S., et al. 2004. Critical role for protein phosphatase 2A heterotrimers in mammalian cell survival. *J. Biol. Chem.* 279: 47732-47739.

MONOS
Satisfaction
Guaranteed

Try **striatin (6): sc-136084**, our highly recommended monoclonal alternative to striatin (N-20).