

# Pendrin (E-20): sc-16894

## BACKGROUND

Pendred syndrome (PDS), an autosomal recessive disorder, is the most common form of syndromic deafness characterized by congenital sensorineural hearing loss and goiter. The gene associated with PDS is mapped to chromosome 7 and encodes a putative transmembrane protein designated Pendrin. Several mutations in the gene have been identified and account for about 10% of hereditary deafness. Pendrin transcripts are expressed at significant levels in the thyroid, inner ear, fetal cochlea and kidney, but expression is drastically reduced in thyroid carcinomas. Pendrin functions as a transporter of chloride and iodide, but not sulfate, in these tissues. It is an apical anion transporter in intercalated cells of proximal tubule and cortical collecting ducts, which mediate renal bicarbonate secretion and  $\text{Cl}^-/\text{OH}^-$ ,  $\text{Cl}^-/\text{HCO}_3^-$  and  $\text{Cl}^-/\text{formate}$  exchange in kidney. Pendrin is expressed throughout the endo-lymphatic duct and sac in distinct areas of the utricle and saccule and in the external sulcus region within the cochlea, where it plays a role in the development of ion gradients.

## REFERENCES

1. Everett, L.A., et al. 1997. Pendred syndrome is caused by mutations in a putative sulphate transporter gene (PDS). *Nat. Genet.* 17: 411-422.
2. Coyle, B., et al. 1998. Molecular analysis of the PDS gene in Pendred syndrome. *Hum. Mol. Genet.* 7: 1105-1112.
3. Everett, L.A., et al. 1999. Expression pattern of the mouse ortholog of the Pendred's syndrome gene (PDS) suggests a key role for Pendrin in the inner ear. *Proc. Natl. Acad. Sci. USA* 96: 9727-9732.
4. Scott, D.A., et al. 1999. The Pendred syndrome gene encodes a chloride-iodide transport protein. *Nat. Genet.* 21: 440-443.

## CHROMOSOMAL LOCATION

Genetic locus: SLC26A4 (human) mapping to 7q22.3; Slc26a4 (mouse) mapping to 12 A3.

## SOURCE

Pendrin (E-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Pendrin of human origin.

## PRODUCT

Each vial contains 200  $\mu\text{g}$  IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-16894 P, (100  $\mu\text{g}$  peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

Pendrin (E-20) is recommended for detection of Pendrin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu\text{g}$  per 100-500  $\mu\text{g}$  of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Pendrin (E-20) is also recommended for detection of Pendrin in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Pendrin siRNA (h): sc-44009, Pendrin siRNA (m): sc-44391, Pendrin shRNA Plasmid (h): sc-44009-SH, Pendrin shRNA Plasmid (m): sc-44391-SH, Pendrin shRNA (h) Lentiviral Particles: sc-44009-V and Pendrin shRNA (m) Lentiviral Particles: sc-44391-V.

Molecular Weight of non-glycosylated Pendrin: 85 kDa.


Molecular Weight of glycosylated Pendrin: 110-140 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



Pendrin (E-20): sc-16894. Western blot analysis of Pendrin expression in Hep G2 whole cell lysate. Note presence of non-denatured and denatured forms.



Pendrin (E-20): sc-16894. Immunofluorescence staining of formalin-fixed Hep G2 cells showing membrane localization.

## SELECT PRODUCT CITATIONS

1. Adler, L., et al. 2008. Molecular mechanisms of epithelial cell-specific expression and regulation of the human anion exchanger (Pendrin) gene. *Am. J. Physiol., Cell Physiol.* 294: C1261-C1276.
2. Udo, K., et al. 2010. Adipose tissue explants and MDCK cells reciprocally regulate their morphogenesis in coculture. *Kidney Int.* 78: 60-68.