

p-Tau (Ser 46): sc-16896

BACKGROUND

Tau, also known as MAPT (microtubule-associated protein tau), MAPTL, MTBT1 or TAU, is a 758 amino acid protein that localizes to the cytoplasm, as well as to the cytoskeleton and the cell membrane, and contains 4 Tau/MAP repeats. Expressed in neuronal tissue and existing as multiple alternatively spliced isoforms, Tau functions to promote microtubule assembly and stability and is thought to be involved in the maintenance of neuronal polarity. Tau may also link microtubules with neural plasma membrane components and, in addition to its role in microtubule stability, is also necessary for cytoskeletal plasticity. Tau is highly subject to a variety of post-translational modifications, including phosphorylation on serine and threonine residues, polyubiquitination (and subsequent proteasomal degradation) and glycation of specific Tau isoforms. Defects in the gene encoding Tau are associated with Alzheimers disease, pallido-ponto-nigral degeneration (PPND), corticobasal degeneration (CBD) and progressive supranuclear palsy (PSP).

REFERENCES

- Hoshi, M., et al. 1996. Regulation of mitochondrial pyruvate dehydrogenase activity by Tau protein kinase I/ glycogen synthase kinase 3 in brain. *Proc. Natl. Acad. Sci. USA* 93: 2719-2723.
- Singh, T.J., et al. 1996. Differential phosphorylation of human Tau isoforms containing three repeats by several protein kinases. *Arch. Biochem. Biophys.* 328: 43-50.
- Tashiro, K., et al. 1997. Somatodendritic localization of phosphorylated Tau in neonatal and adult rat cerebral cortex. *Neuroreport* 8: 2797-2801.
- Iqbal, K., et al. 1998. Mechanisms of neurofibrillary degeneration and the formation of neurofibrillary tangles. *J. Neural. Transm. Suppl.* 53: 169-180.
- Zhong, J., et al. 1999. Hyperphosphorylated Tau in SY5Y cells: similarities and dissimilarities to abnormally hyperphosphorylated Tau from Alzheimer disease brain. *FEBS Lett.* 453: 224-228.
- Iqbal, K., et al. 2000. Mechanism of neurofibrillary degeneration and pharmacologic therapeutic approach. *J. Neural. Transm. Suppl.* 59: 213-222.
- Hashiguchi, M., et al. 2000. 14-3-3 ζ is an effector of Tau protein phosphorylation. *J. Biol. Chem.* 275: 25247-25254.
- Taniguchi, T., et al. 2001. Phosphorylation of Tau is regulated by PKN. *J. Biol. Chem.* 276: 10025-10031.
- Alonso, A., et al. 2001. Hyperphosphorylation induces self-assembly of Tau into tangles of paired helical filaments/straight filaments. *Proc. Natl. Acad. Sci. USA* 98: 6923-6928.

CHROMOSOMAL LOCATION

Genetic locus: MAPT (human) mapping to 17q21.31.

SOURCE

p-Tau (Ser 46) is available as either goat (sc-16896) or rabbit (sc-16896-R) polyclonal affinity purified antibody raised against a short amino acid sequence containing Ser 46 phosphorylated Tau of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-16896 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

p-Tau (Ser 46) is recommended for detection of Ser 46 phosphorylated Tau of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

p-Tau (Ser 46) is also recommended for detection of correspondingly phosphorylated Tau in additional species, including equine, canine and bovine.

Suitable for use as control antibody for Tau siRNA (h): sc-36614, Tau shRNA Plasmid (h): sc-36614-SH and Tau shRNA (h) Lentiviral Particles: sc-36614-V.

Molecular Weight of p-Tau: 46-80 kDa.

Positive Controls: SK-N-SH cell lysate: sc-2410.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.