

SCYL2 (S-14): sc-169263

BACKGROUND

SCYL2 (SCY1-like 2), also known as CVAK104 (coated vesicle-associated kinase of 104 kDa), is a 929 amino acid protein that contains one HEAT repeat and one protein kinase domain. Localized to a variety of locations within the cell, including the perinuclear region of the cytoplasm, the trans-golgi network membrane and clathrin-coated vesicles, SCYL2 exists as a component of AP2-containing Clathrin coated structures. SCYL2 is thought to function as a Ser/Thr protein kinase, exhibiting kinase activity towards adaptor complex proteins, possibly regulating Clathrin-dependent trafficking events between the golgi network and endosomes. Additionally, SCYL2 may play a role in the sorting of SNARE proteins and, ultimately, in embryonic development.

REFERENCES

- Hanks, S.K., Quinn, A.M. and Hunter, T. 1988. The protein kinase family: conserved features and deduced phylogeny of the catalytic domains. *Science* 241: 42-52.
- Nagase, T., Kikuno, R., Ishikawa, K.I., Hirose, M. and Ohara, O. 2000. Prediction of the coding sequences of unidentified human genes. XVI. The complete sequences of 150 new cDNA clones from brain which code for large proteins *in vitro*. *DNA Res.* 7: 65-73.
- Conner, S.D. and Schmid, S.L. 2005. CVAK104 is a novel poly-L-lysine-stimulated kinase that targets the β 2-subunit of AP2. *J. Biol. Chem.* 280: 21539-21544.
- Düwel, M. and Ungewickell, E.J. 2006. Clathrin-dependent association of CVAK104 with endosomes and the *trans*-Golgi network. *Mol. Biol. Cell* 17: 4513-4525.
- Schmid, E.M., Ford, M.G., Burtey, A., Praefcke, G.J., Peak-Chew, S.Y., Mills, I.G., Benmerah, A. and McMahon, H.T. 2006. Role of the AP2 β -appendage hub in recruiting partners for clathrin-coated vesicle assembly. *PLoS Biol.* 4: e262.
- Borner, G.H., Rana, A.A., Forster, R., Harbour, M., Smith, J.C. and Robinson, M.S. 2007. CVAK104 is a novel regulator of clathrin-mediated SNARE sorting. *Traffic* 8: 893-903.

CHROMOSOMAL LOCATION

Genetic locus: SCYL2 (human) mapping to 12q23.1; Scyl2 (mouse) mapping to 10 C2.

SOURCE

SCYL2 (S-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of SCYL2 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-169263 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SCYL2 (S-14) is recommended for detection of SCYL2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other SCYL family members.

SCYL2 (S-14) is also recommended for detection of SCYL2 in additional species, including equine, bovine and avian.

Suitable for use as control antibody for SCYL2 siRNA (h): sc-95795, SCYL2 siRNA (m): sc-153279, SCYL2 shRNA Plasmid (h): sc-95795-SH, SCYL2 shRNA Plasmid (m): sc-153279-SH, SCYL2 shRNA (h) Lentiviral Particles: sc-95795-V and SCYL2 shRNA (m) Lentiviral Particles: sc-153279-V.

Molecular Weight of SCYL2: 104 kDa.

Positive Controls: T-47D cell lysate: sc-2293 or mouse brain extract: sc-2253.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.