

T2R60 (C-11): sc-169502

BACKGROUND

T2R60 (taste receptor type 2 member 60), also known as TAS2R60 or T2R56 (taste receptor type 2 member 56), is a 318 amino acid multi-pass membrane protein that belongs to the G-protein coupled receptor T2R family. T2R60 acts as a receptor that may play a role in the perception of bitterness, and is also thought to be involved in sensing the chemical composition of gastrointestinal content. As a gustducin-linked receptor, the activity of T2R60 may stimulate G_{α_i} (α gustducin), mediate PLC β 2 activation and lead to the gating of TRPM5. While expressed in subsets of taste receptor cells of the tongue and palate epithelium, T2R60 is found exclusively in gustducin-positive cells. The gene that encodes T2R60 contains 957 bases and maps to human chromosome 7q35. Chromosome 7 houses over 1,000 genes, comprises nearly 5% of the human genome and has been linked to osteogenesis imperfecta, Pendred syndrome, lissencephaly, citrullinemia and Shwachman-Diamond syndrome.

REFERENCES

- Iwasaki, S., Iwasaki, S., Usami, S., Abe, S., Isoda, H., Watanabe, T. and Hoshino, T. 2001. Long-term audiological feature in Pendred syndrome caused by PDS mutation. *Arch. Otolaryngol. Head Neck Surg.* 127: 705-708.
- Montmayeur, J.P. and Matsunami, H. 2002. Receptors for bitter and sweet taste. *Curr. Opin. Neurobiol.* 12: 366-371.
- Conte, C., Ebeling, M., Marcuz, A., Nef, P. and Andres-Barquin, P.J. 2002. Identification and characterization of human taste receptor genes belonging to the TAS2R family. *Cytogenet. Genome Res.* 98: 45-53.
- Margolskee, R.F. 2002. Molecular mechanisms of bitter and sweet taste transduction. *J. Biol. Chem.* 277: 1-4.
- Zhang, Y., Hoon, M.A., Chandrashekar, J., Mueller, K.L., Cook, B., Wu, D., Zuker, C.S. and Ryba, N.J. 2003. Coding of sweet, bitter, and umami tastes: different receptor cells sharing similar signaling pathways. *Cell* 112: 293-301.
- Shi, P., Zhang, J., Yang, H. and Zhang, Y.P. 2003. Adaptive diversification of bitter taste receptor genes in mammalian evolution. *Mol. Biol. Evol.* 20: 805-814.
- Go, Y., Satta, Y., Takenaka, O. and Takahata, N. 2005. Lineage-specific loss of function of bitter taste receptor genes in humans and nonhuman primates. *Genetics* 170: 313-326.
- Fischer, A., Gilad, Y., Man, O. and Pääbo, S. 2005. Evolution of bitter taste receptors in humans and apes. *Mol. Biol. Evol.* 22: 432-436.
- Reiner, D., Sapoznik, S. and Sapir, T. 2006. Lissencephaly 1 linking to multiple diseases: mental retardation, neurodegeneration, schizophrenia, male sterility, and more. *Neuromolecular Med.* 8: 547-565.

CHROMOSOMAL LOCATION

Genetic locus: TAS2R60 (human) mapping to 7q35.

SOURCE

T2R60 (C-11) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a C-terminal cytoplasmic domain of T2R60 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-169502 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

T2R60 (C-11) is recommended for detection of T2R60 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other T2R family members.

Suitable for use as control antibody for T2R60 siRNA (h): sc-89617, T2R60 shRNA Plasmid (h): sc-89617-SH and T2R60 shRNA (h) Lentiviral Particles: sc-89617-V.

Molecular Weight of T2R60: 36 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.