

# WDR92 (C-15): sc-169808

## BACKGROUND

WD-repeats are motifs that are found in a variety of proteins and are characterized by a conserved core of 40-60 amino acids that commonly form a tertiary propeller structure. While proteins that contain WD-repeats participate in a wide range of cellular functions, they are generally involved in regulatory mechanisms concerning chromatin assembly, cell cycle control, signal transduction, RNA processing, apoptosis and vesicular trafficking. WDR92 (WD-repeat-containing protein 92), also known as WD repeat-containing protein Monad, is a 357 amino acid protein that contains 6 WD-repeats. Expressed in a variety of tissues, with highest levels present in testis, WDR92 is suggested to influence apoptosis. The gene encoding WDR92 maps to human chromosome 2, which houses over 1,400 genes and comprises nearly 8% of the human genome and comprises nearly 8% of the human genome. A number of genetic diseases are linked to genes on chromosome 2 including Harlequin ichthyosis, sitosterolemia and Alström syndrome.

## REFERENCES

1. Van der Voorn, L. and Ploegh, H.L. 1992. The WD-40 repeat. *FEBS Lett.* 307: 131-134.
2. Neer, E.J., Schmidt, C.J., Nambudripad, R. and Smith, T.F. 1994. The ancient regulatory-protein family of WD-repeat proteins. *Nature* 371: 297-300.
3. Smith, T.F., Gaitatzes, C., Saxena, K. and Neer, E.J. 1999. The WD repeat: a common architecture for diverse functions. *Trends Biochem. Sci.* 24: 181-185.
4. Zumsteg, U., Muller and P.Y., Miserez, A.R. 2000. Alstrom syndrome: confirmation of linkage to chromosome 2p12-13 and phenotypic heterogeneity in three affected sibs. *J. Med. Genet.* 37: E8.
5. Shulenin, S., Schriml, L.M., Remaley, A.T., Fojo, S., Brewer, B., Allikmets and R., Dean, M. 2001. An ATP-binding cassette gene (ABCG5) from the ABCG (White) gene subfamily maps to human chromosome 2p21 in the region of the Sitosterolemia locus. *Cytogenet. Cell Genet.* 92: 204-208.

## CHROMOSOMAL LOCATION

Genetic locus: WDR92 (human) mapping to 2p14; Wdr92 (mouse) mapping to 11 A2.

## SOURCE

WDR92 (C-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of WDR92 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-169808 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

WDR92 (C-15) is recommended for detection of WDR92 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other WDR family members.

WDR92 (C-15) is also recommended for detection of WDR92 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for WDR92 siRNA (h): sc-94404, WDR92 siRNA (m): sc-155329, WDR92 shRNA Plasmid (h): sc-94404-SH, WDR92 shRNA Plasmid (m): sc-155329-SH, WDR92 shRNA (h) Lentiviral Particles: sc-94404-V and WDR92 shRNA (m) Lentiviral Particles: sc-155329-V.


Molecular Weight of WDR92: 40 kDa.

Positive Controls: Sol8 nuclear extract: sc-2157 or mouse testis extract: sc-2405.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



WDR92 (C-15): sc-169808. Western blot analysis of WDR92 expression in Sol8 nuclear extract (A) and mouse testis tissue extract (B).

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.