



p-IL-3R β (Ser 585): sc-17008

BACKGROUND

The human IL-3, IL-5 and GM-CSF receptors are each composed of both unique α subunits and a common 130 kDa β subunit. The α subunits are low affinity ligand binding proteins while the β subunits do not themselves bind ligand, but are required for high affinity binding by the α subunits. In contrast, the mouse IL-3 receptor has two distinct β subunits, one that functions only in IL-3 mediated cell signaling and a second that is shared with IL-5 and GM-CSF. The murine β -subunits are 91% homologous at the amino acid level but only 56% homologous to the human β subunit. Although neither the murine nor the human β subunit contains tyrosine kinase domains, both activate tyrosine phosphorylation mediated signaling pathways.

REFERENCES

- Hayashida, K., Kitamura, R., Gorman, D.M., Yokata, T. and Miyajima, A. 1990. Molecular cloning of a second subunit of the receptor for human granulocyte-macrophage colony-stimulating factor (GM-CSF): reconstitution of a high affinity GM-CSF receptor. *Proc. Natl. Acad. Sci. USA* 87: 9655-9659.
- Tavernier, J., Devos, R., Cornelis, S., Tuyperis, T., VanderHeydon, J., Qiers, W. and Platenok, G. 1992. A human high affinity interleukin-5 receptor (IL-5R) is composed of an IL-5 specific chain and a β chain shared with the receptor for GM-CSF. *Cell* 66: 1175-1184.
- Hara, T. and Miyajima, A. 1992. Two distinct functional receptors for mouse interleukin-3. *EMBO J.* 11: 1875-1884.
- Sakamaki, K., Miyajima, I., Kitamura, T. and Miyajima, A. 1992. Critical cytoplasmic domains of the common β subunit of the human GM-CSF, IL-3 and IL-5 receptors for growth signal transduction and tyrosine phosphorylation. *EMBO J.* 11: 3541-3549.
- Park, L.S., Martin, U., Sorenson, R., Luhr, S., Morrissey, P.J., Cosman, D. and Larsen, A. 1992. Cloning of the low-affinity murine granulocyte-macrophage colony-stimulating factor receptor and reconstitution of a high-affinity receptor complex. *Proc. Natl. Acad. Sci. USA* 89: 4295-4299.
- Miyajima, A., Kitamura, T., Harada, N., Yokota, T. and Arai, K. 1992. Cytokine receptors and signal transduction. *Ann. Rev. Immunol.* 10: 295-331.
- Goodall, G.J., Bagley, C.J., Vadas, M.A. and Lopez, A.F. 1993. A model for the interaction of the GM-CSF, IL-3 and IL-5 receptors with their ligands. *Growth Fact.* 8: 87-97.
- Rao, P., Kitamura, T., Miyajima, A. and Mufson, R.A. 1995. Human IL-3 receptor signaling: rapid induction of phosphatidylcholine hydrolysis is independent of protein kinase C but dependent on tyrosine phosphorylation in transfected NIH/3T3 cells. *J. Immunol.* 154: 1664-1674.
- Guthridge, M.A., Stomski, F.C., Barry, E.F., Winnall, W., Woodcock, J.M., McClure, B.J., Dottore, M., Berndt, M.C. and Lopez, A.F. 2000. Site-specific Serine phosphorylation of the IL-3 receptor is required for hemopoietic cell survival. *Mol. Cell* 6: 99-108.

CHROMOSOMAL LOCATION

Genetic locus: CSF2RB (human) mapping to 22q13.1; Csf2rb1 (mouse) mapping to 15 E1.

SOURCE

p-IL-3R β (Ser 585) is available as either goat (sc-17008) or rabbit (sc-17008-R) polyclonal affinity purified antibody raised against a short amino acid sequence containing phosphorylated Ser 585 of IL-3R β of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-17008 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

p-IL-3R β (Ser 585) is recommended for detection of Ser 585 phosphorylated IL-3R β of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for IL-3R β siRNA (h): sc-40060.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent) and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE** Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.