

Tim13/AB (C-20): sc-17065

BACKGROUND

The majority of mitochondrial-directed proteins are encoded by the nuclear genome and are transported to the mitochondria via regulated processes involving the mitochondrial Tom and Tim proteins. The mitochondrial Tim protein family is comprised of a large group of evolutionarily conserved proteins that are found in most eukaryotes. Import of nuclear-encoded precursor proteins into and across the mitochondrial inner membrane is mediated by two distinct complexes, the Tim23 complex and the Tim22 complex, which differ in their substrate specificity. Defects in Tim proteins are implicated in several neuro-degenerative diseases, suggesting important roles for Tim proteins in development and health. Tim13A and Tim13B, which map to human chromosomes 19p13.3-p13.1 and 19p13.3, respectively, are highly similar to yeast Tim13, which localizes to the yeast mitochondrial intermembrane space. Tim13 forms a complex with Tim8, which interacts with the amino-terminal domain of Tim23.

REFERENCES

- Jin, H., Kendall, E., Freeman, T.C., Roberts, R.G. and Vetrie, D.L. 1999. The human family of Deafness/Dystonia peptide (DDP) related mitochondrial import proteins. *Genomics* 61: 259-267.
- Bauer, M.F., Rothbauer, U., Muhlenbein, N., Smith, R.J., Gerbitz, K., Neupert, W., Brunner, M. and Hofmann, S. 1999. The mitochondrial TIM22 preprotein translocase is highly conserved throughout the eukaryotic kingdom. *FEBS Lett.* 464: 41-47.
- Rassow, J., Dekker, P.J., van Wilpe, S., Meijer, M. and Soll, J. 1999. The preprotein translocase of the mitochondrial inner membrane: function and evolution. *J. Mol. Biol.* 286: 105-120.
- Koehler, C. M., Leuenerberger, D., Merchant, S., Renold, A., Junne, T. and Schatz, G. 1999. Human deafness dystonia syndrome is a mitochondrial disease. *Proc. Natl. Acad. Sci. USA* 96: 2141-216.
- Paschen, S.A., Rothbauer, U., Kaldi, K., Bauer, M.F., Neupert, W. and Brunner, M. 2000. The role of the TIM8-13 complex in the import of Tim23 into mitochondria. *EMBO J.* 19: 6392-6400.
- Bauer, M.F. and Neupert, W. 2001. Import of proteins into mitochondria: a novel pathomechanism for progressive neurodegeneration. *J. Inher. Metab. Dis.* 24: 166-180.
- LocusLink Report (LocusID: 26518). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: TIMM13 (human) mapping to 19p13.3; Timm13 (mouse) mapping to 10 C1.

SOURCE

Tim13A/B (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of Tim13A of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-17065 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Tim13/AB (C-20) is recommended for detection of Tim13A and Tim13B of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Tim13/AB (C-20) is also recommended for detection of Tim13A and Tim13B in additional species, including equine, bovine and porcine.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.