

p-p53 (Ser 315)-R: sc-17106-R

BACKGROUND

p53 is a DNA-binding, oligomerization domain- and transcription activation domain-containing tumor suppressor that upregulates growth arrest and apoptosis-related genes in response to stress signals, thereby influencing programmed cell death, cell differentiation and cell cycle control mechanisms. p53 localizes to the nucleus yet can be chaperoned to the cytoplasm by the negative regulator MDM2, an E3 ubiquitin ligase that is upregulated in the presence of active p53, where MDM2 polyubiquitinates p53 for proteasome targeting. p53 can assemble into tetramers in the absence of DNA, fluctuates between latent and active (DNA-binding) conformations, and is differentially activated through posttranslational modifications including phosphorylation and acetylation. Mutations in the DNA-binding domain (DBD) (amino acids 110-286) of p53 can compromise energetically favorable association with *cis* elements and are implicated in several human cancers. Phosphorylation of p53 at residue Thr 155 is mediated by the COP9 signalosome (CSN) and targets p53 to ubiquitin-26S Proteasome-dependent degradation.

REFERENCES

- Hupp, T.R., Meek, D.W., Midgley, C.A. and Lane, D.P. 1992. Regulation of the specific DNA binding function of p53. *Cell* 71: 875-886.
- Levine, A. J. 1997. p53, the cellular gatekeeper for growth and division. *Cell* 88: 323-331.
- Ashcroft, M. and Vousden, K.H. 1999. Regulation of p53 stability. *Oncogene* 18: 7637-7643.
- Soussi, T., Dehouche, K. and Beroud, C. 2000. p53 website and analysis of p53 gene mutations in human cancer: forging a link between epidemiology and carcinogenesis. *Hum. Mutat.* 15: 105-113.
- Chene, P. 2001. The role of tetramerization in p53 function. *Oncogene* 20: 2611-2617.
- Minamoto, T., Buschmann, T., Habelhah, H., Matusевич, E., Tahara, H., Boerresen-Dale, A.L., Harris, C., Sidransky, D. and Ronai, Z. 2001. Distinct pattern of p53 phosphorylation in human tumors. *Oncogene* 20: 3341-3347.
- Bech-Otschir, D., Kraft, R., Huang, X., Henklein, P., Kapelari, B., Pollmann, C. and Dubiel, W. 2001. COP9 signalosome-specific phosphorylation targets p53 to degradation by the ubiquitin system. *EMBO J.* 20: 1630-1639.
- LocusLink Report (LocusID: 7157). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: TP53 (human) mapping to 17p13.1.

SOURCE

p-p53 (Ser 315)-R is a rabbit polyclonal antibody raised against a short amino acid sequence containing Ser 315 phosphorylated p53 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-17106 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

p-p53 (Ser 315)-R is recommended for detection of Ser 315 phosphorylated p53 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

p-p53 (Ser 315)-R is also recommended for detection of correspondingly phosphorylated p53 in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for p53 siRNA (h): sc-29435, p53 shRNA Plasmid (h): sc-29435-SH, p53 shRNA and p53 shRNA (h) Lentiviral Particles: sc-29435-V.

Molecular Weight of p-p53: 53 kDa.

Positive Controls: MCF7+ etoposide cell lysate: sc-2281.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent), Western Blotting Luminol Reagent: sc-2048 and Lambda Phosphatase: sc-200312A. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
 Satisfation
 Guaranteed

Try **p-p53 (59.Ser 315): sc-135772** or **p-p53 (64.Ser 315): sc-135773**, our highly recommended monoclonal alternatives to p-p53 (Ser 315).