

TAZ (G-20): sc-17131

BACKGROUND

The transcriptional co-activator with PDZ-binding motif (TAZ) is a 14-3-3-binding molecule. The highly conserved and ubiquitously expressed 14-3-3 proteins regulate differentiation, cell cycle progression and apoptosis by binding intracellular phosphoproteins involved in signal transduction. TAZ may link events at the plasma membrane and cytoskeleton to nuclear transcription in a manner that can be regulated by 14-3-3. TAZ shares homology with the WW domain of Yes-associated protein (YAP) and functions as a transcriptional co-activator by binding to the PPXY motif present on transcription factors. TAZ recognizes immunoreactive protein bands in lysates from MDCK, NIH-3T3 and 293T cells. In addition, COS7, Hep G2, CHO and HeLa cells express endogenous TAZ. 14-3-3 binding requires TAZ phosphorylation on a single Serine 89 residue, resulting in the inhibition of TAZ transcriptional co-activation through 14-3-3-mediated nuclear export.

CHROMOSOMAL LOCATION

Genetic locus: WWTR1 (human) mapping to 3q25.1; Wwtr1 (mouse) mapping to 3 D.

SOURCE

TAZ (G-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of TAZ of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-17131 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

TAZ (G-20) is recommended for detection of TAZ of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

TAZ (G-20) is also recommended for detection of TAZ in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for TAZ siRNA (h): sc-38568, TAZ siRNA (m): sc-38569, TAZ shRNA Plasmid (h): sc-38568-SH, TAZ shRNA Plasmid (m): sc-38569-SH, TAZ shRNA (h) Lentiviral Particles: sc-38568-V and TAZ shRNA (m) Lentiviral Particles: sc-38569-V.


Molecular Weight of TAZ: 45 kDa.

Positive Controls: TAZ (h): 293T Lysate: sc-170455 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



TAZ (G-20): sc-17131. Western blot analysis of TAZ expression in non-transfected: sc-117752 (A) and human TAZ transfected: sc-170455 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **TAZ (1F1): sc-293183**, our highly recommended monoclonal alternative to TAZ (G-20).