

# YAP (S-20): sc-17140

## BACKGROUND

The Yes-associated protein, otherwise known as YAP, is a 14-3-3 binding molecule that was originally recognized by virtue of its ability to bind to the SH3 domain of Yes. The binding of YAP to 14-3-3 requires the phosphorylation of a homologous serine residue (Ser 112) in the YAP 14-3-3 binding motif. The highly conserved and ubiquitously expressed 14-3-3 proteins regulate differentiation, cell cycle progression and apoptosis by binding intracellular phosphoproteins involved in signal transduction. YAP, which has a molecular mass of approximately 65 kDa, may link events at the plasma membrane and cytoskeleton to inhibition of transcription in the nucleus in a manner regulated by 14-3-3 proteins. YAP shares homology with the WW domain of TAZ, transcriptional co-activator with PDZ-binding motif, which functions as a transcriptional co-activator by binding to the PPXY motif present in transcription factors. YAP is expressed at high levels in the lung, placenta, prostate, ovary and testis.

## CHROMOSOMAL LOCATION

Genetic locus: YAP1 (human) mapping to 11q13.1; Yap1 (mouse) mapping to 9 A1.

## SOURCE

YAP (S-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of YAP of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-17140 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-17140 X, 200 µg/0.1 ml.

## APPLICATIONS

YAP (S-20) is recommended for detection of YAP of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

YAP (S-20) is also recommended for detection of YAP in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for YAP siRNA (h): sc-38637, YAP siRNA (m): sc-38638, YAP shRNA Plasmid (h): sc-38637-SH, YAP shRNA Plasmid (m): sc-38638-SH, YAP shRNA (h) Lentiviral Particles: sc-38637-V and YAP shRNA (m) Lentiviral Particles: sc-38638-V.

YAP (S-20) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of YAP: 65 kDa.

Positive Controls: PC-3 cell lysate: sc-2220 or HeLa nuclear extract: sc-2120.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

## DATA



YAP (S-20): sc-17140. Immunoperoxidase staining of formalin fixed, paraffin-embedded human placenta tissue showing nuclear and cytoplasmic staining of decidual cells.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **YAP (G-6): sc-376830** or **YAP (H-9): sc-271134**, our highly recommended monoclonal alternatives to YAP (S-20). Also, for AC, HRP, FITC, PE, Alexa Fluor<sup>®</sup> 488 and Alexa Fluor<sup>®</sup> 647 conjugates, see **YAP (G-6): sc-376830**.