

Sox-30 (S-20): sc-17373

BACKGROUND

Sox-30 (SRY (sex determining region Y)-box 30) encodes a member of the Sox (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. Sox-30 is expressed in testis and may act as a transcriptional regulator after forming a protein complex with other proteins. Sox-30 may be involved in the differentiation of developing male germ cells. Two transcript variants encoding distinct isoforms have been identified for the human Sox-30 gene. Sox family transcription factors influence cell differentiation, development and sex determination. Sox-30 contains a unique DNA binding domain, known as the high mobility group (HMG) box, that is related to that of the testis determining gene, SRY. The highly complex group of Sox genes cluster at a minimum of 40 different loci that rapidly diverged in various animal lineages. Several Sox genes have been identified, and members of this family have been shown to be conserved during evolution and to play key roles during animal development.

REFERENCES

- Laudet, V., Stehelin, D. and Clevers, H. 1993. Ancestry and diversity of the HMG box superfamily. *Nucleic Acids Res.* 21: 2493-2501.
- Kuhlbrodt, K., Herbarth, B., Sock, E., Hermans-Borgmeyer, I. and Wegner, M. 1998. Sox-10, a novel transcriptional modulator in glial cells. *J. Neurosci.* 18: 237-250.
- Arsic, N., Rajic, T., Stanojic, S., Goodfellow, P.N. and Stevanovic, M. 1998. Characterisation and mapping of the human Sox-14 gene. *Cytogenet. Cell Genet.* 83: 139-146.
- Osaki, E., Nishina, Y., Inazawa, J., Copeland, N.G., Gilbert, D.J., Jenkins, N.A., Ohsugi, M., Tezuka, T., Yoshida, M. and Semba, K. 1999. Identification of a novel SRY-related gene and its germ cell-specific expression. *Nucleic Acids Res.* 27: 2503-2510.
- Sasai, Y. 2001. Roles of Sox factors in neural determination: conserved signaling in evolution? *Int. J. Dev. Biol.* 45: 321-326.

CHROMOSOMAL LOCATION

Genetic locus: SOX30 (human) mapping to 5q33.3.

SOURCE

Sox-30 (S-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Sox-30 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-17373 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Sox-30 (S-20) is recommended for detection of Sox-30 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Sox-30 (S-20) is also recommended for detection of Sox-30 in additional species, including canine and porcine.

Suitable for use as control antibody for Sox-30 siRNA (h): sc-36529, Sox-30 shRNA Plasmid (h): sc-36529-SH and Sox-30 shRNA (h) Lentiviral Particles: sc-36529-V.

Sox-30 (S-20) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of Sox-30: 94 kDa.

Positive Controls: SK-N-MC nuclear extract: sc-2154, IMR-32 nuclear extract: sc-2148 or A-431 nuclear extract: sc-2122.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.