

# MYPT1 (N-15): sc-17433

## BACKGROUND

Myosin phosphatase target subunit 1 (MYPT1), also called Myosin-binding subunit of Myosin phosphatase, is one of the subunits and an integral component of the Myosin phosphatase. Myosin phosphatase regulates the interaction of Actin and Myosin downstream of the guanosine triphosphatase Rho, which inhibits Myosin phosphatase through the action of Rho-kinase. MYPT1 promotor contains one Sp1 transcription factor binding site, suggesting that MYPT1 is a housekeeping gene. Myotonic dystrophy protein kinase phosphorylates MYPT1 at Tyrosine 654 to regulate Myosin II phosphorylation. Inhibition of Myosin light chain phosphatase results in  $Ca^{2+}$  sensitization of smooth muscle contraction. This inhibition is modulated through phosphorylation of MYPT1 by a ZIP-like kinase, which associates with MYPT1 and phosphorylates the inhibitory site in smooth muscle. The phosphorylation of MYPT1 by protein kinase C results in altered dephosphorylation of Myosin by attenuating the binding of protein phosphatase 1 catalytic subunit (PP1c) and the phosphorylated Myosin light chain to MYPT1. PP1c interacts with at least four binding sites on the amino-terminus of MYPT1. MYPT2, a novel isoform of MYPT1, also interacts with PP1c. MYPT1 is localized on stress fibers; it is distributed close to the cell membrane and at cell-cell contacts to regulate Myosin phosphatase activity.

## CHROMOSOMAL LOCATION

Genetic locus: PPP1R12A (human) mapping to 12q15-q21; Ppp1r12a (mouse) mapping to 10 C.

## SOURCE

MYPT1 (N-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of MYPT1 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-17433 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

MYPT1 (N-15) is recommended for detection of MYPT1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000). MYPT1 (N-15) is also recommended for detection of MYPT1 in additional species, including canine, bovine, porcine and avian.

Suitable for use as control antibody for MYPT1 siRNA (h): sc-37240, MYPT1 siRNA (m): sc-37241, MYPT1 shRNA Plasmid (h): sc-37240-SH, MYPT1 shRNA Plasmid (m): sc-37241-SH, MYPT1 shRNA (h) Lentiviral Particles: sc-37240-V and MYPT1 shRNA (m) Lentiviral Particles: sc-37241-V.


Molecular Weight of MYPT1: 130 kDa.

Positive Controls: DU 145 cell lysate: sc-2268, 3T3-L1 cell lysate: sc-2243 or MYPT1 (h2): 293T Lysate: sc-114575.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



MYPT1 (N-15): sc-17433. Western blot analysis of MYPT1 expression in non-transfected 293T: sc-117752 (A), human MYPT1 transfected 293T: sc-114575 (B) and 3T3-L1 (C) whole cell lysates.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **MYPT1 (C-6): sc-514261**, our highly recommended monoclonal alternative to MYPT1 (N-15). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **MYPT1 (C-6): sc-514261**.