

Myosin Va (N-20): sc-17706

BACKGROUND

Myosin Va, a member of the unconventional Myosin family, is a non-muscle Myosin that structurally combines elements from both nonmuscle Myosin type I and nonmuscle Myosin type II. Class V unconventional myosins, which include Myosin Va and Myosin Vb, are nonfilamentous, Actin-binding enzymes that appear to be expressed ubiquitously. Myosin V proteins are regulated by their heavy chain phosphorylation, which occurs in the carboxy-terminal tail domain. The mouse Myosin Va gene is also known as the mouse dilute gene because mutations in this gene cause the coat color in mice to lighten. Mutations in the Myosin Va gene also result in the onset of severe neurological defects shortly after birth. Defects in the human Myosin Va gene, which maps to chromosome 15q21.2, have been implicated in Griscelli disease, a rare autosomal recessive disorder characterized by pigmentary dilution, variable cellular immunodeficiency and onset of acute, uncontrolled lymphocyte and macrophage activation.

CHROMOSOMAL LOCATION

Genetic locus: MYO5A (human) mapping to 15q21.2; Myo5a (mouse) mapping to 9 D.

SOURCE

Myosin Va (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Myosin Va of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-17706 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Myosin Va (N-20) is recommended for detection of Myosin Va of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Myosin Va (N-20) is also recommended for detection of Myosin Va in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for Myosin Va siRNA (h): sc-35995, Myosin Va siRNA (m): sc-35996, Myosin Va shRNA Plasmid (h): sc-35995-SH, Myosin Va shRNA Plasmid (m): sc-35996-SH, Myosin Va shRNA (h) Lentiviral Particles: sc-35995-V and Myosin Va shRNA (m) Lentiviral Particles: sc-35996-V.


Molecular Weight of Myosin Va: 190 kDa.

Positive Controls: A-375 cell lysate: sc-3811, HeLa whole cell lysate: sc-2200 or KNRK whole cell lysate: sc-2214.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Myosin Va (N-20): sc-17706. Western blot analysis of Myosin Va expression in A-375 whole cell lysate.



Myosin Va (N-20): sc-17706. Immunofluorescence staining of methanol-fixed KNRK cells showing cytoplasmic localization.

SELECT PRODUCT CITATIONS

- Passeron, T., et al. 2004. Cyclic AMP promotes a peripheral distribution of melanosomes and stimulates melanophilin/Slac2-a and Actin association. *FASEB J.* 18: 989-991.
- Ogiwara, K. and Hata, K. 2009. Melanoma cell differentiation induced by lupeol separates into two stages: morphological and functional changes. *J. Nat. Med.* 63: 323-326.
- Yoshii, A., et al. 2013. A Myosin Va mutant mouse with disruptions in glutamate synaptic development and mature plasticity in visual cortex. *J. Neurosci.* 33: 8472-8482.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.


 MONOS
Satisfaction
Guaranteed

Try **Myosin Va (G-4): sc-365986**, our highly recommended monoclonal alternative to Myosin Va (N-20).