## SANTA CRUZ BIOTECHNOLOGY, INC.

# CtBP (E-12): sc-17759



#### BACKGROUND

CtBP1 is a cellular phosphoprotein that associates with various proteins and functions as a corepressor of transcription. CtBP1 and the related protein CtBP2 are characterized as C-terminal binding protein of adenovirus E1A, and they preferentially associate with the E1A via a 5-amino acid motif, PLDLS, to repress E1A induced oncogenesis and cellular transformation. CtBP1 is expressed from embryo to adult, but CtBP2 is mainly expressed during embryogenesis. During skeletal and T cell development, CtBP1 and CtBP2 associate with the PLDLSL domain of  $\delta$ EF1, a cellular zinc finger-homeodomain protein, and thereby enhances  $\delta$ EF1 induced transcriptional silencing. In addition, CtBP complexes with CtIP, a protein that recognizes distinctly different protein motifs from CtBP. CtIP binds to the BRCT repeats within the breast cancer gene BRCA1 and enables CtBP to influence BRCA1 activity. CtIP/CtBP binding to BRCA1 inhibits the transactivation of the p21 promoter, and it is critical for regulating p21 transcription in response to DNA damage.

# **CHROMOSOMAL LOCATION**

Genetic locus: CTBP1 (human) mapping to 4p16.3, CTBP2 (human) mapping to 10q26.13; Ctbp1 (mouse) mapping to 5 B1, Ctbp2 (mouse) mapping to 7 F3.

## SOURCE

CtBP (E-12) is a mouse monoclonal antibody raised against amino acids 1-440 of CtBP1 of human origin.

### PRODUCT

Each vial contains 200  $\mu$ g lgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CtBP (E-12) is available conjugated to agarose (sc-17759 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-17759 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-17759 PE), fluorescein (sc-17759 FITC), Alexa Fluor<sup>®</sup> 488 (sc-17759 AF488), Alexa Fluor<sup>®</sup> 546 (sc-17759 AF546), Alexa Fluor<sup>®</sup> 594 (sc-17759 AF594) or Alexa Fluor<sup>®</sup> 647 (sc-17759 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor<sup>®</sup> 680 (sc-17759 AF680) or Alexa Fluor<sup>®</sup> 790 (sc-17759 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

#### **APPLICATIONS**

CtBP (E-12) is recommended for detection of CtBP1 and CtBP2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:200-1:2000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of CtBP: 48 kDa.

Positive Controls: Neuro-2A whole cell lysate: sc-364185, NRK whole cell lysate: sc-364197 or SH-SY5Y cell lysate: sc-3812.

#### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### DATA



CtBP (E-12): sc-17759. Western blot analysis of CtBP expression in SH-SY5Y (A), Neuro-2A (B), Sol8 (C), 3T3-L1 (D), C6 (E) and NRK (F) whole cell lysates.



CtBP (E-12): sc-17759. Immunofluorescence detection of CtBP in formalin-fixed HeLa cells showing nuclear localization. Detection reagent used: m-IgG\kappa BP-PE: sc-516141 (**A**). Immunoperoxidase staining of formalin fixed, paraffin-embedded human small intestine tissue showing nuclear staining of glandular cells (**B**).

#### SELECT PRODUCT CITATIONS

- 1. Deltour, S., et al. 2002. The human candidate tumor suppressor gene HIC1 recruits CtBP through a degenerate GLDLSKK motif. Mol. Cell. Biol. 22: 4890-4901.
- Yang, L., et al. 2011. ncRNA- and Pc2 methylation-dependent gene relocation between nuclear structures mediates gene activation programs. Cell 147: 773-788.
- 3. Siles, L., et al. 2013. ZEB1 imposes a temporary stage-dependent inhibition of muscle gene expression and differentiation via CtBP-mediated transcriptional repression. Mol. Cell. Biol. 33: 1368-1382.
- Malik, M.Q., et al. 2014. Small ubiquitin-like modifier (SUMO)-mediated repression of the *Xenopus* oocyte 5 S rRNA genes. J. Biol. Chem. 289: 35468-35481.
- Sánchez-Tilló, E., et al. 2015. ZEB1 and TCF4 reciprocally modulate their transcriptional activities to regulate Wnt target gene expression. Oncogene 34: 5760-5770.
- Janeckova, L., et al. 2016. Wnt signaling inhibition deprives small intestinal stem cells of clonogenic capacity. Genesis 54: 101-114.
- 7. Usui, N., et al. 2017. Foxp1 regulation of neonatal vocalizations via cortical development. Genes Dev. 31: 2039-2055.
- 8. Hrckulak, D., et al. 2018. Wnt Effector TCF4 is dispensable for Wnt signaling in human cancer cells. Genes 9: 439.
- 9. Zou, T., et al. 2019. Organoid-derived C-Kit+/SSEA4-human retinal progenitor cells promote a protective retinal microenvironment during transplantation in rodents. Nat. Commun. 10: 1205.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.