

# DNA pol $\delta$ cat (C-2): sc-17777

## BACKGROUND

DNA replication, recombination and repair, all of which are necessary for genome stability, require the presence of exonucleases. In DNA replication, these enzymes are involved in the processing of Okazaki fragments, whereas in DNA repair, they function to excise damaged DNA fragments and correct recombinational mismatches. Exonucleases involved in these processes include DNA polymerases, including DNA pol  $\delta$  and  $\epsilon$ . DNA pol  $\delta$  consists of two subunits, p125 which interacts directly with the sliding DNA clamp protein PCNA, and p50. DNA pol  $\delta$  can be regulated by cell cycle proteins. DNA pol  $\epsilon$  is a multiple subunit enzyme, the catalytic subunit of which is encoded by the POL2 gene. The exact reactions catalyzed by DNA pol  $\delta$  and  $\epsilon$  on leading and lagging strands have not yet been elucidated.

## REFERENCES

- Lee, M.Y., et al. 1984. Further studies on calf thymus DNA polymerase  $\delta$  purified to homogeneity by a new procedure. *Biochemistry* 23: 1906-1913.
- Hamatake, R.K., et al. 1990. Purification and characterization of DNA polymerase II from the yeast *Saccharomyces cerevisiae*. Identification of the catalytic core and a possible holoenzyme form of the enzyme. *J. Biol. Chem.* 265: 4072-4083.

## CHROMOSOMAL LOCATION

Genetic locus: POLD1 (human) mapping to 19q13.33; Pold1 (mouse) mapping to 7 B4.

## SOURCE

DNA pol  $\delta$  cat (C-2) is a mouse monoclonal antibody raised against amino acids 1-300 of DNA pol  $\delta$  cat of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>2b</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

DNA pol  $\delta$  cat (C-2) is recommended for detection of DNA pol  $\delta$  cat of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for DNA pol  $\delta$  cat siRNA (h): sc-37777, DNA pol  $\delta$  cat siRNA (m): sc-37778, DNA pol  $\delta$  cat shRNA Plasmid (h): sc-37777-SH, DNA pol  $\delta$  cat shRNA Plasmid (m): sc-37778-SH, DNA pol  $\delta$  cat shRNA (h) Lentiviral Particles: sc-37777-V and DNA pol  $\delta$  cat shRNA (m) Lentiviral Particles: sc-37778-V.


Molecular Weight of DNA pol  $\delta$  cat: 125 kDa.

Positive Controls: MDA-MB-231 cell lysate: sc-2232, HeLa whole cell lysate: sc-2200 or Jurkat whole cell lysate: sc-2204.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



DNA pol  $\delta$  cat (C-2): sc-17777. Western blot analysis of DNA pol  $\delta$  cat expression in Jurkat (A), HL-60 (B), MCF7 (C), HeLa (D), MDA-MB-231 (E) and RAW 264.7 (F) whole cell lysates. Detection reagent used: m-IgG<sub>2b</sub> BP-HRP: sc-542741.



DNA pol  $\delta$  cat (C-2): sc-17777. Western blot analysis of DNA pol  $\delta$  cat expression in HeLa (A), MDA-MB-231 (B), Ramos (C), BYDP (D) and WEHI-231 (E) whole cell lysates.

## SELECT PRODUCT CITATIONS

- Stanislav, N., et al. 2005. Proliferating cell nuclear antigen (PCNA) may function as a double homotrimer complex in the mammalian cell. *J. Biol. Chem.* 280: 13888-13894.
- Takaya, J., et al. 2013. Protein interaction and cellular localization of human CDC45. *J. Biochem.* 153: 381-388.
- Maskey, R.S., et al. 2017. Spartan deficiency causes accumulation of Topoisomerase 1 cleavage complexes and tumorigenesis. *Nucleic Acids Res.* 45: 4564-4576.
- Layer, J.V., et al. 2020. Polymerase  $\delta$  promotes chromosomal rearrangements and imprecise double-strand break repair. *Proc. Natl. Acad. Sci. USA* 117: 27566-27577.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.