

# TBX5 (H-20): sc-17865

## BACKGROUND

The T-box (TBX) motif is present in a family of genes whose structural features and expression patterns support their involvement in developmental gene regulation. The TBX gene family are largely conserved throughout metazoan evolution, and these genes code for putative transcription factors that share a uniquely defining DNA-binding domain. TBX genes are a family of developmental regulators with more than 20 members recently identified in invertebrates and vertebrates. Mutations in TBX genes are associated with the onset of several human diseases. Our understanding of functional mechanisms of TBX products has come mainly from the prototypical T/Brachyury, which is a transcription activator. The TBX genes constitute a family of transcriptional regulatory genes that are implicated in a variety of developmental processes ranging from the formation of germ layers to the organizational patterning of the central nervous system.

## REFERENCES

1. Law, D.J., et al. 1995. Identification, characterization, and localization to chromosome 17q21-22 of the human TBX2 homolog, member of a conserved developmental gene family. *Mamm. Genome* 6: 793-797.
2. Agulnik, S.I., et al. 1998. Cloning, mapping, and expression analysis of TBX15, a new member of the T-Box gene family. *Genomics* 51: 68-75.
3. Dheen, T., et al. 1999. Zebrafish TBX-c functions during formation of mid-line structures. *Development* 126: 2703-2713.
4. He, M.I., et al. 1999. Transcription repression by *Xenopus* ET and its human ortholog TBX3, a gene involved in ulnar-mammary syndrome. *Proc. Natl. Acad. Sci. USA* 96: 10212-10217.
5. Begemann, G. and Ingham, P.W. 2000. Developmental regulation of TBX5 in zebrafish embryogenesis. *Mech. Dev.* 90: 299-304.
6. Ahn, D.G., et al. 2000. TBX20, a new vertebrate T-box gene expressed in the cranial motor neurons and developing cardiovascular structures in zebrafish. *Mech. Dev.* 95: 253-258.

## CHROMOSOMAL LOCATION

Genetic locus: TBX5 (human) mapping to 12q24.1; Tbx5 (mouse) mapping to 5 F.

## SOURCE

TBX5 (H-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of TBX5 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-17865 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

TBX5 (H-20) is recommended for detection of TBX5 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

TBX5 (H-20) is also recommended for detection of TBX5 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for TBX5 siRNA (h): sc-37020, TBX5 siRNA (m): sc-37021, TBX5 shRNA Plasmid (h): sc-37020-SH, TBX5 shRNA Plasmid (m): sc-37021-SH, TBX5 shRNA (h) Lentiviral Particles: sc-37020-V and TBX5 shRNA (m) Lentiviral Particles: sc-37021-V.


Molecular Weight of TBX5: 57 kDa.

Positive Controls: HeLa nuclear extract: sc-2120.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



TBX5 (H-20): sc-17865. Western blot analysis of TBX5 expression in HeLa nuclear extract.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

**MONOS**  
Satisfaction  
Guaranteed

Try **TBX5 (A-6): sc-515536** or **TBX5 (A-4): sc-376952**, our highly recommended monoclonal alternatives to TBX5 (H-20).