
## SANTA CRUZ BIOTECHNOLOGY, INC.

# TBX1 (G-17): sc-17874



## BACKGROUND

The T-box (TBX) motif is present in a family of genes whose structural features and expression patterns support their involvement in developmental gene regulation. The TBX gene family are largely conserved throughout metazoan evolution, and these genes code for putative transcription factors that share a uniquely defining DNA-binding domain. TBX genes are a family of developmental regulators with more than 20 members recently identified in invertebrates and vertebrates. Mutations in TBX genes are associated with the onset of several human diseases. Our understanding of functional mechanisms of TBX products has come mainly from the prototypical T/Brachyury, which is a transcription activator. The TBX genes constitute a family of transcriptional regulatory genes that are implicated in a variety of developmental processes ranging from the formation of germ layers to the organizational patterning of the central nervous system.

## REFERENCES

- Law, D.J., Gebuhr, T., Garvey, N., Agulnik, S.I. and Silver, L.M. 1995. Identification, characterization, and localization to chromosome 17q21-22 of the human TBX2 homolog, member of a conserved developmental gene family. Mamm. Genome 6: 793-797.
- Agulnik, S.I., Papaioannou, V.E. and Silver, L.M. 1998. Cloning, mapping, and expression analysis of TBX15, a new member of the T-box gene family. Genomics 51: 68-75.
- Dheen, T., Sleptsova-Friedrich, I., Xu, Y., Clark, M., Lehrach, H., Gong, Z. and Korzh, V. 1999. Zebrafish TBX-C functions during formation of midline structures. Development 126: 2703-2713.
- He, M.I.., Wen, L., Campbell, C.E., Wu, J.Y. and Rao, Y. 1999. Transcription repression by Xenopus ET and its human ortholog TBX3, a gene involved in ulnar-mammary syndrome. Proc. Natl. Acad. Sci. USA 96: 10212-10217.
- Ahn, D.G., Ruvinsky, I., Oates, A.C., Silver, L.M. and Ho, R.K. 2000. TBX20, a new vertebrate T-box gene expressed in the cranial motor neurons and developing cardiovascular structures in zebrafish. Mech. Dev. 95: 253-258.
- Begemann, G. and Ingham, P.W. 2000. Developmental regulation of TBX5 in zebrafish embryogenesis. Mech. Dev. 90: 299-304.

## CHROMOSOMAL LOCATION

Genetic locus: TBX1 (human) mapping to 22q11; Tbx1 (mouse) mapping to 16 11.35 cM.

#### SOURCE

TBX1 (G-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of TBX1 of human origin.

## **STORAGE**

Store at 4° C, \*\*D0 NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### PRODUCT

Each vial contains 200  $\mu g$  lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-17874 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-17874 X, 200  $\mu$ g/0.1 ml.

## **APPLICATIONS**

TBX1 (G-17) is recommended for detection of TBX1 of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TBX1 siRNA (h): sc-38467.

TBX1 (G-17) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

## **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluores-cence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941.

#### **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.