



## COL4A3 (N-16): sc-18177

### BACKGROUND

The extensive family of COL gene products (collagens) is composed of several chain types, including fibril-forming interstitial Collagens (Types I, II, III and V) and basement membrane Collagens (Type IV), each type containing multiple isoforms. Collagens are fibrous, extracellular matrix proteins with high tensile strength and are the major components of connective tissue, such as tendons and cartilage. All collagens contain a triple helix domain and frequently show lateral self-association in order to form complex connective tissues. Several collagens also play a role in cell adhesion, important for maintaining normal tissue architecture and function.

### REFERENCES

1. Bateman, J.F., et al. 1996. In Comper, W.D., ed. Extracellular Matrix. Amsterdam: Harwood. 2: 22-67.
2. McCarthy, J.B., et al. 1996. Cell adhesion to collagenous matrices. Biopolymers 40: 371-381.
3. Engel, J. 1997. Versatile collagens in invertebrates. Science 277: 1785-1786.
4. Cremer, M.A., et al. 1998. The cartilage collagens: a review of their structure, organization, and role in the pathogenesis of experimental arthritis in animals and in human rheumatic disease. J. Mol. Med. 76: 275-288.
5. Boskey, A.L., et al. 1999. Collagen and bone strength. J. Bone Miner. Res. 14: 330-335.
6. Alberio, L. and Dale, G.L. 1999. Platelet-collagen interactions: membrane receptors and intracellular signaling pathways. Eur. J. Clin. Invest. 29: 1066-1076.
7. Kalluri, R., 2002. Discovery of type IV Collagen non-collagenous domains as novel integrin ligands and endogenous inhibitors of angiogenesis. Cold Spring Harb. Symp. Quant. Biol. 67: 255-266.
8. Pescucci, C., et al. 2003. Type-IV Collagen related diseases. J. Nephrol. 16: 314-316.
9. Kim, S.J., et al. 2004. Effects of type IV Collagen and laminin on the cryopreservation of human embryonic stem cells. Stem Cells 22: 950-961.

### CHROMOSOMAL LOCATION

Genetic locus: COL4A3 (human) mapping to 2q36.3

### SOURCE

COL4A3 (N-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Collagen  $\alpha 3$  Type IV of human origin.

### PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-18177 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

COL4A3 (N-16) is recommended for detection of Collagen  $\alpha 3$  Type IV of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for COL4A3 siRNA (h): sc-105228, COL4A3 shRNA Plasmid (h): sc-105228-SH and COL4A3 shRNA (h) Lentiviral Particles: sc-105228-V.

Molecular Weight of COL4A3: 160-190 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.