

Hepsin (V-16): sc-18572

BACKGROUND

Extracellular proteases mediate the digestion of neighboring extracellular matrix components in initial tumor growth, allow desquamation of tumor cells into the surrounding environment, provide the basis for invasion of basement membranes in targeted metastatic organs and are required for release and activation of many growth and angiogenic factors. Hepsin (also known as TMPRSS1) is a type II transmembrane serine protease in mammalian cells that is highly expressed on the surface of hepatocytes. Hepsin is frequently overexpressed in several tumors, suggesting that it is a candidate protease in the invasive process and growth capacity of tumor cells. The basal promoter region of the Hepsin gene contains potential binding sites for SP1, AP2, C/EBP, LF-A1 and E box, which may be responsible for the ubiquitous expression of the protein, which shows preferential expression in liver and kidney. Hepsin is located at the plasma membrane, with its catalytic subunit (C-terminal half) at the cell surface and its N-terminus facing the cytosol. Hepsin has been shown to play a role in normal cell growth, embryogenesis, hepatocyte growth, blood coagulation and fertilization. In addition, Hepsin converts zymogen Factor VII to Factor VIIa, which is capable of initiating a coagulation pathway on the cell surface and ultimately leads to Thrombin formation.

CHROMOSOMAL LOCATION

Genetic locus: HPN (human) mapping to 19q13.12.

SOURCE

Hepsin (V-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Hepsin of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-18572 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Hepsin (V-16) is recommended for detection of Hepsin of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Hepsin (V-16) is also recommended for detection of Hepsin in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for Hepsin siRNA (h): sc-60782; and as shRNA Plasmid control antibody for Hepsin shRNA Plasmid (h): sc-60782-SH.


Molecular Weight of Hepsin: 51 kDa.

Positive Controls: Hepsin (h): 293T Lysate: sc-114224 or Hep G2 cell lysate: sc-2227.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Hepsin (V-16): sc-18572. Western blot analysis of Hepsin expression in non-transfected 293T: sc-117752 (A), human Hepsin transfected 293T: sc-114224 (B) and Hep G2 (C) whole cell lysates.

SELECT PRODUCT CITATIONS

- Xing, P., et al. 2011. Clinical and biological significance of hepsin overexpression in breast cancer. *J. Investig. Med.* 59: 803-810.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **Hepsin (2D5): sc-517056**, our highly recommended monoclonal alternative to Hepsin (V-16).