

LRP (C-14): sc-18703

BACKGROUND

Tumor cells that are insensitive to anticancer drugs often have a multidrug-resistant (MDR) phenotype. Proteins associated with this phenomenon are transport-associated proteins such as P-glycoprotein, multidrug resistance protein 1, lung resistance-related protein (LRP) and breast cancer resistance protein (BCRP). The LRP protein, which is identified as the major vault protein (MVP), is overexpressed in various multidrug-resistant cancer cell lines and clinical samples. The promoter of LRP is TATA-less and contains an inverted CCAAT-box and a Sp1 site located near a p53 binding motif. LRP has two alternative splice variants, which differ from each other within the 5'-leader. The long-LRP isoform is ubiquitously expressed and represents an almost constant portion of the total LRP mRNA in many different normal tissues. LRP is the major component of the multimeric ribonucleoprotein complexes, with several copies of an untranslated RNA, which has been shown to transport along cytoskeletal-based cellular tracks. In conclusion, LRP protein mediates drug resistance, perhaps via a transport process.

REFERENCES

1. Scheffer, G.L., et al. 1995. The drug resistance-related protein LRP is the human major vault protein. *Nat. Med.* 1: 578-582.
2. Herrmann, C., et al. 1999. Recombinant major vault protein is targeted to neuritic tips of PC12 cells. *J. Cell Biol.* 144: 1163-1172.
3. Scheffer, G.L., et al. 2000. Lung resistance-related protein/major vault protein and vaults in multidrug-resistant cancer. *Curr. Opin. Oncol.* 12: 550-556.
4. Lange, C., et al. 2000. Cloning and initial analysis of the human multidrug resistance-related MVP/LRP gene promoter. *Biochem. Biophys. Res. Commun.* 278: 125-133.

CHROMOSOMAL LOCATION

Genetic locus: MVP (human) mapping to 16p11.2.

SOURCE

LRP (C-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of drug resistance-related protein of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-18703 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

LRP (C-14) is recommended for detection of drug resistance-related protein of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for LRP siRNA (h): sc-35824, LRP shRNA Plasmid (h): sc-35824-SH and LRP shRNA (h) Lentiviral Particles: sc-35824-V.


Molecular Weight of LRP: 110 kDa.

Positive Controls: Caki-1 cell lysate: sc-2224, T98G cell lysate: sc-2294 or U-87 MG cell lysate: sc-2411.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



LRP (C-14): sc-18703. Immunoperoxidase staining of formalin fixed, paraffin-embedded human nasopharynx tissue showing cytoplasmic and nuclear staining of respiratory epithelial cells.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **LRP (1014): sc-23916** or **LRP (E-2): sc-390134**, our highly recommended monoclonal alternatives to LRP (C-14).