

# ATR (N-19): sc-1887

## BACKGROUND

Members of the PIK (phosphatidylinositol kinase)-related kinase family are high molecular weight kinases involved in cell cycle progression, DNA recombination and detection of DNA damage. One member of the PI 3-/PI 4-kinase family is ATR (ataxia-telangiectasia- and Rad3-related), also known as FRP1 (for FRAP-related protein 1). ATR is most closely related to ATM, a protein kinase encoded by the gene mutated in ataxia telangiectasia. ATR is also closely related to three of the family members involved in checkpoint function: Mei-41 (*Drosophila*), Mec1p (*S. cerevisiae*) and Rad3 (*Schizosaccharomyces pombe*), and as such may be the functional human counterpart of these proteins. This kinase has been shown to phosphorylate checkpoint kinase CHK1, checkpoint proteins Rad17 and Rad9, as well as tumor suppressor protein BRCA1. In addition, ATR is essential for early embryonic development. The protein encoded by the human ATR gene localizes to intranuclear foci after DNA damage or inhibition of replication.

## CHROMOSOMAL LOCATION

Genetic locus: ATR (human) mapping to 3q23; Atr (mouse) mapping to 9 E3.3.

## SOURCE

ATR (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of ATR of human origin.

## PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-1887 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

ATR (N-19) is recommended for detection of ATR of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ATR (N-19) is also recommended for detection of ATR in additional species, including canine and porcine.

Suitable for use as control antibody for ATR siRNA (h): sc-29763, ATR siRNA (m): sc-29764, ATR shRNA Plasmid (h): sc-29763-SH, ATR shRNA Plasmid (m): sc-29764-SH, ATR shRNA (h) Lentiviral Particles: sc-29763-V and ATR shRNA (m) Lentiviral Particles: sc-29764-V.

Molecular Weight of ATR: 250 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, A-431 whole cell lysate: sc-2201 or K-562 whole cell lysate: sc-2203.


## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## DATA



ATR (N-19): sc-1887. Western blot analysis of ATR expression in HeLa (A), A-431 (B) and K-562 (C) whole cell lysates.



ATR (N-19): sc-1887. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear and cytoplasmic localization.

## SELECT PRODUCT CITATIONS

- Hu, B., et al. 2001. The radioresistance to killing of A1-5 cells derives from activation of the Chk1 pathway. *J. Biol. Chem.* 276: 17693-17698.
- Mu, X.F., et al. 2011. DNA damage-sensing kinases mediate the mouse 2-cell embryo's response to genotoxic stress. *Biol. Reprod.* 85: 524-535.
- Harrigan, J.A., et al. 2011. Replication stress induces 53BP1-containing OPT domains in G<sub>1</sub> cells. *J. Cell Biol.* 193: 97-108.
- Hu, C., et al. 2012. Roles of kruppel-associated box (KRAB)-associated co-repressor KAP1 Ser-473 phosphorylation in DNA damage response. *J. Biol. Chem.* 287: 18937-18952.
- Page, J., et al. 2012. Inactivation or non-reactivation: what accounts better for the silence of sex chromosomes during mammalian male meiosis? *Chromosoma* 121: 307-326.
- Pérez-Castro, A.J., et al. 2012. Rad9B responds to nucleolar stress through ATR and JNK signalling, and delays the G<sub>1</sub>-S transition. *J. Cell Sci.* 125: 1152-1164.
- Liu, G., et al. 2012. Replication fork stalling and checkpoint activation by a PKD1 locus mirror repeat polypurine-polypyrimidine (Pu-Py) tract. *J. Biol. Chem.* 287: 33412-33423.
- Ray, A., et al. 2013. NER initiation factors, DDB2 and XPC, regulate UV radiation response by recruiting ATR and ATM kinases to DNA damage sites. *DNA Repair* 12: 273-283.
- Yuan, F., et al. 2013. Overexpressed DNA polymerase  $\epsilon$  regulated by JNK/c-Jun contributes to hypermutagenesis in bladder cancer. *PLoS ONE* 8: e69317.

**MONOS**  
Satisfaction  
Guaranteed

Try **ATR (C-1): sc-515173**, our highly recommended monoclonal alternative to ATR (N-19). Also, for AC, HRP, FITC, PE, Alexa Fluor<sup>®</sup> 488 and Alexa Fluor<sup>®</sup> 647 conjugates, see **ATR (C-1): sc-515173**.