

# HAI-2 (L-20): sc-18936

## BACKGROUND

Tissue damage, such as hepatic and renal injury, initiates hepatocyte growth factor activator (HGFAC)-mediated limited proteolytic activation of the inactive single-chain precursor form of HGF. Initially, HGFAC is produced as a precursor protein, which is activated by limited proteolysis and is neutralized by specific inhibitors known as HGF activator inhibitors, designated HAIs. HAIs belong to the Kunitz-type serine protease inhibitor family. HAIs target HGF activator and are involved in the regulation of proteolytic activation of HGF in injured tissues. Human HAI-1 transcript is expressed in various human tissues, such as adult placenta, kidney, pancreas, prostate and small intestine, and fetal kidney and lung. It translates into a 478 amino acid protein. The human HAI-2 gene maps to chromosome 19q13.1 and encodes a 252 amino acid protein, also designated human placental bikunin or kop (Kunitz domain containing protein over-expressed in pancreatic cancer). HAI-1 and HAI-2 are produced in membrane-associated forms, which are secreted as active, proteolytically truncated proteins.

## REFERENCES

1. Shimomura, T., et al. 1997. Hepatocyte growth factor activator inhibitor, a novel Kunitz-type serine protease inhibitor. *J. Biol. Chem.* 272: 6370-6376.
2. Marlor, C.W., et al. 1997. Identification and cloning of human placental bikunin, a novel serine protease inhibitor containing two Kunitz domains. *J. Biol. Chem.* 272: 12202-12208.
3. Kawaguchi, T., et al. 1997. Purification and cloning of hepatocyte growth factor activator inhibitor type 2, a Kunitz-type serine protease inhibitor. *J. Biol. Chem.* 272: 27558-27564.
4. Muller-Pillasch, F., et al. 1998. Cloning of a new Kunitz-type protease inhibitor with a putative transmembrane domain overexpressed in pancreatic cancer. *Biochim. Biophys. Acta* 1395: 88-95.
5. Online Mendelian Inheritance in Man, OMIM™. 2000. Johns Hopkins University, Baltimore, MD. MIM Number: 605123. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## CHROMOSOMAL LOCATION

Genetic locus: SPINT2 (human) mapping to 19q13.1; Spint2 (mouse) mapping to 7 A3.

## SOURCE

HAI-2 (L-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of HAI-2 of mouse origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-18936 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

HAI-2 (L-20) is recommended for detection of precursor and mature HAI-2 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for HAI-2 siRNA (m): sc-39557, HAI-2 shRNA Plasmid (m): sc-39557-SH and HAI-2 shRNA (m) Lentiviral Particles: sc-39557-V.

Molecular Weight of HAI-2: 23-31 kDa.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



HAI-2 (L-20): sc-18936. Western blot analysis of HAI-2 expression in non-transfected: sc-117752 (A) and mouse HAI-2 transfected: sc-126933 (B) 293T whole cell lysates.



HAI-2 (L-20): sc-18936. Western blot analysis of His-tagged mouse recombinant HAI-2.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **HAI-2 (H-9): sc-398119**, our highly recommended monoclonal alternative to HAI-2 (L-20).