

# Tctex1 (N-19): sc-19184

## BACKGROUND

Dyneins are multisubunit, high molecular weight ATPases that interact with microtubules to generate force by converting the chemical energy of ATP into the mechanical energy of movement. Cytoplasmic or axonemal Dynein heavy, intermediate, light and light-intermediate chains are all components of minus end-directed motors; the complex transports cellular cargos towards the central region of the cell. Axonemal Dynein motors contain one to three non-identical heavy chains and cause a sliding of microtubules in the axonemes of cilia and flagella in a mechanism necessary for cilia to beat and propel the cell. Cytoplasmic Dynein is an approximately 12 subunit complex of two heavy chains, two intermediate chains to anchor Dynein to its cargo, four smaller intermediate chains and several light chains. It performs functions necessary for cell survival such as organelle transport and centrosome assembly. The carboxy-terminus of Dynein is important for microtubule-dependent motility and is highly conserved, while the amino-terminal regions are more variable. Tctex1 is a cytoplasmic dynein light chain found in a complex with Na<sup>+</sup> CP type X $\alpha$  (SCN10A). Tctex1, also designated CW-1 or TCTEL1 is expressed in heart, placenta, skeletal muscle, kidney, pancreas, spleen, prostate, testis, ovary, ileum and colon. Several proteins regulate Dynein activity, including dynectin, LIS1 and NudEL(NudE-like).

## REFERENCES

1. Watanabe, T.K., et al. 1996. Cloning, expression, and mapping of TCTEL1, a putative human homologue of murine Tcte1, to 6q. *Cytogenet. Cell Genet.* 73: 153-156.
2. Asai, D.J., et al. 2004. The dynein heavy chain family. *J. Eukaryot. Microbiol.* 51: 23-29.
3. Chuang, J.Z., et al. 2005. The dynein light chain Tctex-1 has a dynein-independent role in actin remodeling during neurite outgrowth. *Dev. Cell* 9: 75-86.
4. Li, J., et al. 2005. NudEL targets dynein to microtubule ends through LIS1. *Nat. Cell Biol.* 7: 686-690.
5. Seetharam, R.N., et al. 2005. High speed sliding of axonemal microtubules produced by outer arm dynein. *Cell Motil. Cytoskeleton* 60: 96-103.
6. He, Y., et al. 2005. Role of cytoplasmic dynein in the axonal transport of microtubules and neurofilaments. *J. Cell Biol.* 168: 697-703.
7. Pfister, K.K., et al. 2005. Cytoplasmic dynein nomenclature. *J. Cell Biol.* 171: 411-413.

## CHROMOSOMAL LOCATION

Genetic locus: DYNLT1 (human) mapping to 6q25.3; DYNLT1a/DYNLT1b/DYNLT1c/DYNLT1f (mouse) mapping to 17 A1.

## SOURCE

Tctex1 (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of Tctex1 of human origin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-19184 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

Tctex1 (N-19) is recommended for detection of Tctex1 of human origin, DYNLT1a, DYNLT1b, DYNLT1c and DYNLT1f of mouse origin, and the corresponding rat homolog by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Tctex1 (N-19) is also recommended for detection of Tctex1 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for Tctex1 siRNA (h): sc-43319, Tctex1 shRNA Plasmid (h): sc-43319-SH and Tctex1 shRNA (h) Lentiviral Particles: sc-43319-V.

Molecular Weight of Tctex1: 14 kDa.

Positive Controls: Y79 cell lysate: sc-2240, IMR-32 cell lysate: sc-2409 or PC-3 cell lysate: sc-2220.

## DATA



Tctex1 (N-19): sc-19184. Western blot analysis of Tctex1 expression in PC-3 (A) and Y79 (B) whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **Tctex1 (H-11): sc-365567**, our highly recommended monoclonal alternative to Tctex1 (N-19).