

11 β -HSD2 (T-12): sc-19262

BACKGROUND

Glucocorticoid hormone action in target tissues is modulated by 11 β -hydroxysteroid dehydrogenase (11 β -HSD), which catalyzes the interconversion of hormonally active C11-hydroxylated corticosteroids (cortisol, corticosterone) and their inactive C11-keto metabolites (cortisone, 11-dehydrocorticosterone). At least two isoforms of 11 β -HSD exist: a low-affinity NADP-dependent dehydrogenase/oxoreductase (11 β -HSD1) and a high-affinity NAD-dependent dehydrogenase (11 β -HSD2). The glycosylated 11 β -HSD1 protein activates cortisol from cortisone, which is widely expressed in mammals, and is most highly expressed in the liver. 11 β -HSD2 inactivates cortisol to cortisone and is expressed in placenta, aldosterone target tissues (kidney, parotid, colon and skin) and pancreas. 11 β -HSD1 may play a role in glucose homeostasis and pathogenesis of a number of disorders including Insulin resistance and obesity. 11 β -HSD2 associates with differentiation or maturation in human colonic epithelia and may serve as a marker in development and disease. In addition, 11 β -HSD2 plays a crucial role in modulating mineralcorticoid and glucocorticoid receptor occupancy by glucocorticoids.

REFERENCES

- Tannin, G.M., et al. 1991. The human gene for 11 β -hydroxysteroid dehydrogenase. Structure, tissue distribution, and chromosomal localization. *J. Biol. Chem.* 266: 16653-16658.
- Albiston, A.L., et al. 1994. Cloning and tissue distribution of the human 11 beta-hydroxysteroid dehydrogenase type 2 enzyme. *Mol. Cell Endocrinol.* 105: 11-17.
- Brown, R.W., et al. 1996. Cloning and production of antisera to human placental 11 β -hydroxysteroid dehydrogenase type 2. *Biochem. J.* 313: 1007-1017.
- Takahashi, K., et al. 1998. 11 β -hydroxysteroid dehydrogenase type II in human colon: a new marker of fetal development and differentiation in neoplasms. *Anticancer Res.* 18: 3381-3388.
- Arcuri, F., et al. 1999. Expression of 11 β -hydroxysteroid dehydrogenase in early pregnancy: implications in human trophoblast-endometrial interactions. *Semin. Reprod. Endocrinol.* 17: 53-61.

CHROMOSOMAL LOCATION

Genetic locus: HSD11B2 (human) mapping to 16q22.

SOURCE

11 β -HSD2 (T-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of 11 β -HSD2 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-19262 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

11 β -HSD2 (T-12) is recommended for detection of 11 β -hydroxysteroid dehydrogenase type 2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for 11 β -HSD2 siRNA (h): sc-41379, 11 β -HSD2 shRNA Plasmid (h): sc-41379-SH and 11 β -HSD2 shRNA (h) Lentiviral Particles: sc-41379-V.


Molecular Weight of 11 β -HSD2: 40 kDa.

Positive Controls: 11 β -HSD2 (h3): 293T Lysate: sc-110103, HeLa whole cell lysate: sc-2200 or HCT 116 cell lysate.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



11 β -HSD2 (T-12): sc-19262. Western blot analysis of 11 β -HSD2 expression in non-transfected: sc-117752 (A) and human 11 β -HSD2 transfected: sc-116955 (B) 293T whole cell lysates.



11 β -HSD2 (T-12): sc-19262. Western blot analysis of 11 β -HSD2 expression in non-transfected: sc-117752 (A) and human 11 β -HSD2 transfected: sc-110103 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

- Cheng, H., et al. 2009. Validation of immature adipogenic status and identification of prognostic biomarkers in myxoid liposarcoma using tissue microarrays. *Hum. Pathol.* E-published.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.