

DNase II (N-13): sc-19271

BACKGROUND

Internucleosomal DNA fragmentation following the activation of endonucleases is the common end point of apoptosis. DNase I, a $\text{Ca}^{2+}/\text{Mg}^{2+}$ -dependent endonuclease ubiquitously expressed in mammalian tissues, has been implicated to mediate internucleosomal DNA degradation in human cells undergoing apoptosis. DNase I is highly polymorphic, and at least six alleles of DNase I are known. DNase II, the ubiquitously expressed acidic deoxyribonuclease, acts downstream of caspase activation and may also induce DNA digestion during apoptosis. DNase I cleaves DNA to 5'-phosphodinucleotide and 5'-phospho-oligonucleotide end-products, whereas DNase II cleaves DNA to 3'-phosphomononucleotide and 3'-phospho-oligonucleotide end-products. The mechanism by which DNase II cuts DNA is similar to DNase I, which produces nicks rather than double-strand cuts. DNase II is usually present in cytoplasm of epithelial cells, but it appears concentrated in the nuclei of lens fibers. In contrast, DNase I is always concentrated in nuclei of epithelial and fiber cells.

REFERENCES

1. Torriglia, A., Chaudun, E., Chany-Fournier, F., Jeanny, J.C., Courtois, Y. and Counis, M.F. 1995. Involvement of DNase II in nuclear degeneration during lens cell differentiation. *J. Biol. Chem.* 270: 28579-28585.
2. Yasuda, T., Takeshita, H., Iida, R., Nakajima, T., Hosomi, O., Nakashima, Y. and Kishi, K. 1998. Molecular cloning of the cDNA encoding human deoxyribonuclease II. *J. Biol. Chem.* 273: 2610-2616.
3. Krieser, R.J. and Eastman, A. 1998. The cloning and expression of human deoxyribonuclease II. A possible role in apoptosis. *J. Biol. Chem.* 273: 30909-30914.
4. Baker, K.P., Baron, W.F., Henzel, W.J. and Spencer, S.A. 1998. Molecular cloning and characterization of human and murine DNase II. *Gene* 215: 281-289.
5. Yasuda, T., Takeshita, H., Iida, R., Kogure, S. and Kishi, K. 1999. A new allele, DNASE1*6, of human deoxyribonuclease I polymorphism encodes an Arg to Cys substitution responsible for its instability. *Biochem. Biophys. Res. Commun.* 260: 280-283.
6. Oliveri, M., Daga, A., Cantoni, C., Lunardi, C., Millo, R. and Puccetti, A. 2001. DNase I mediates internucleosomal DNA degradation in human cells undergoing drug-induced apoptosis. *Eur. J. Immunol.* 31: 743-751.

CHROMOSOMAL LOCATION

Genetic locus: DNASE2 (human) mapping to 19q13.2; Dnase2a (mouse) mapping to 8 C3.

SOURCE

DNase II (N-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of DNase II of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-19271 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

DNase II (N-13) is recommended for detection of DNase II of human, rat and, to a lesser extent, mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for DNase II siRNA (h): sc-41507, DNase II shRNA Plasmid (h): sc-41507-SH and DNase II shRNA (h) Lentiviral Particles: sc-41507-V.

Molecular Weight of DNase II: 32 kDa.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



DNase II (N-13): sc-19271. Western blot analysis of DNase II expression in mouse liver extract.



DNase II (N-13): sc-19271. Western blot analysis of DNase II expression in mouse liver tissue extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.