SANTA CRUZ BIOTECHNOLOGY, INC.

ACP1α/β (N-15): sc-19333



BACKGROUND

Regulation of intracellular concentrations of flavoenzymes and flavin coenzymes is essential for proper cell homeostasis. Red cell acid phosphatase, known as ACP1, catalyzes the transfer of phosphate from phosphate ester substrates to suitable acceptor alcohols such as methanol and glycerol. ACP is a genetically polymorphic, cytoplasmic low-molecular-weight flavin mononucleotide phosphatase that regulates the intracellular concentrations of flavin coenzymes. The human ACP1 gene maps to chromosome 2p25 and encodes a pair of isozymes, Bf (alpha) and Bs (beta) The ACP1 alpha and beta isozymes are not glycosylated. Both ACP1-alpha and ACP1beta isozymes are 157 amino acids in length; however the two forms differ in sequence over an internal 34 residue segment. The two isoforms are believed to differ in substrate specificity.

REFERENCES

- Golden, V.L. and Sensabaugh, G.F. 1986. Phenotypic variation in the phosphotransferase activity of human red cell acid phosphatase (ACP1). Hum. Genet. 72: 340-343.
- Dissing, J. and Sensabaugh, G.F. 1987. Human red cell acid phosphatase (ACP1): evidence for differences in the primary structure of the two isozymes encoded by the ACP1*B allele. Biochem. Genet. 25: 919-927.
- 3. Dissing, J., et al. 1991. Human red cell acid phosphatase (ACP1). The amino acid sequence of the two isozymes Bf and Bs encoded by the ACP1*B allele. J. Biol. Chem. 266: 20619-20625.
- 4. Dissing, J. and Johnsen, A.H. 1992. Human red cell acid phosphatase (ACP1): the primary structure of the two pairs of isozymes encoded by the ACP1*A and ACP1*C alleles. Biochim. Biophys. Acta 1121: 261-268.
- 5. Wo, Y.Y., et al. 1992. Sequencing, cloning, and expression of human red cell-type acid phosphatase, a cytoplasmic phosphotyrosyl protein phosphatase. J. Biol. Chem. 267: 10856-10865.

CHROMOSOMAL LOCATION

Genetic locus: ACP1 (human) mapping to 2p25.3; Acp1 (mouse) mapping to 12 A2.

SOURCE

ACP1 α/β (N-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of ACP1 of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-19333 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

ACP1 α/β (N-15) is recommended for detection of ACP1 α and ACP1 β of mouse, rat, human and origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1,000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3,000).

ACP1 α/β (N-15) is also recommended for detection of ACP1 α and ACP1 β in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for ACP1 siRNA (h): sc-108019, ACP1 siRNA (m): sc-44359, ACP1 shRNA Plasmid (h): sc-108019-SH, ACP1 shRNA Plasmid (m): sc-44359-SH, ACP1 shRNA (h) Lentiviral Particles: sc-108019-V and ACP1 shRNA (m) Lentiviral Particles: sc-44359-V.

Positive Controls: Hep G2 cell lysate: sc-2227 or ACP1 (h): 293T Lysate: sc-111661.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2,000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA





ACP1 α/β (N-15): sc-19333. Western blot analysis of ACP1 expression in non-transfected: sc-117752 (**A**) and human ACP1 transfected: sc-111661 (**B**) 293T whole cell lysates.

ACP1 α/β (N-15): sc-19333. Immunofluorescence staining of methanol-fixed KNRK cells showing cytoplasmic localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS Satisfation Guaranteed

Try **ACP1 (D-3):** sc-390190 or **ACP1\alpha/\beta (B-2):** sc-398459, our highly recommended monoclonal alternatives to ACP1 α/β (N-15).