

# ORC6 (V-15): sc-19731

## BACKGROUND

The initiation of DNA replication is a multi-step process that depends on the formation of pre-replication complexes, which trigger initiation. Among the proteins required for establishing these complexes are the origin recognition complex (ORC) proteins. ORC proteins bind specifically to origins of replication where they serve as scaffold for the assembly of additional initiation factors. Human ORC subunits 1-6 are expressed in the nucleus of proliferating cells and tissues, such as the testis. ORC1 and ORC2 are both expressed at equivalent concentrations throughout the cell cycle; however, only ORC2 remains stably bound to chromatin. ORC4 and ORC6 are also expressed constantly throughout the cell cycle. ORC2, ORC3, ORC4 and ORC5 form a core complex upon which the ORC6 and ORC1 assemble. The formation of this core complex suggests ORC proteins play a crucial role in the G<sub>1</sub>-S transition in mammalian cells.

## REFERENCES

1. Quintana, D.G., Hou, Z., Thome, K.C., Hendricks, M., Saha, P. and Dutta, A. 1997. Identification of the HsORC4, a member of the human origin of replication recognition complex. *J. Biol. Chem.* 272: 28247-28251.
2. Mendez, J. and Stillman, B. 2000. Chromatin association of human origin recognition complex, Cdc6, and minichromosome maintenance proteins during the cell cycle: assembly of prereplication complexes in late mitosis. *Mol. Cell. Biol.* 20: 8602-8612.
3. Dhar, S.K. and Dutta, A. 2000. Identification and characterization of the human ORC6 homolog. *J. Biol. Chem.* 275: 34983-34988.
4. Thome, K.C., Dhar, S.K., Quintana, D.G., Delmolino, L., Shahsafaei, A. and Dutta, A. 2000. Subsets of human origin recognition complex (ORC) subunits are expressed in non-proliferating cells and associate with non-ORC proteins. *J. Biol. Chem.* 275: 35233-35241.
5. Kreitz, S., Ritzi, M., Baack, M. and Knippers, R. 2000. The human origin-recognition-complex protein 1 dissociates from chromatin during S phase in HeLa cells. *J. Biol. Chem.* 276: 6337-6342.
6. Natale, D.A., Li, C.J., Sun, W.H. and DePamphilis, M.L. 2000. Selective instability of ORC1 protein accounts for the absence of functional origin recognition complexes during the M-G<sub>1</sub> transition in mammals. *EMBO J.* 19: 2728-2738.
7. Vashee, S., Simancek, P., Challberg, M.D. and Kelly, T.J. 2001. Assembly of the human origin recognition complex. *J. Biol. Chem.* 276: 26666-26673.
8. Dhar, S.K., Delmolino, L. and Dutta, A. 2001. Architecture of the human origin recognition complex. *J. Biol. Chem.* 276: 29067-29071.

## CHROMOSOMAL LOCATION

Genetic locus: ORC6 (human) mapping to 16q11.2; Orc6 (mouse) mapping to 8 C3.

## SOURCE

ORC6 (V-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ORC6 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-19731 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-19731 X, 200 µg/0.1 ml.

## APPLICATIONS

ORC6 (V-15) is recommended for detection of ORC6 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ORC6 (V-15) is also recommended for detection of ORC6 in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for ORC6 siRNA (h): sc-38161, ORC6 siRNA (m): sc-38162, ORC6 shRNA Plasmid (h): sc-38161-SH, ORC6 shRNA Plasmid (m): sc-38162-SH, ORC6 shRNA (h) Lentiviral Particles: sc-38161-V and ORC6 shRNA (m) Lentiviral Particles: sc-38162-V.

ORC6 (V-15) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of ORC6: 30 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, U-2 OS cell lysate: sc-2295 or SW480 cell lysate: sc-2219.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.