SANTA CRUZ BIOTECHNOLOGY, INC.

Fe65 (N-20): sc-19750



BACKGROUND

 γ -secretase cleaves the cell surface protein amyloid protein precursor (APP) at the transmembrane region into an extracelluar peptide (Amyloid- β) and an intracelluar tail fragment. The cytoplasmic tail of APP forms a multimeric complex with Fe65 (also known as APBB1 for APP binding protein family B member 1). Specifically, Fe65 binds the YENPTY sequence in the cytoplasmic tail of APP. Fe65 is a nuclear adaptor protein widely expressed in the brain, including hippocampus and isocortex. In the cell, Fe65 and APP colocalize to the ER and Golgi. The interaction between APP and Fe65 increases the translocation of APP to the cell surface and the subsequent secretion of Amyloid- β . Fe65 and APP localize with Mena, a cell-adhesion protein, and Fe65 regulates APP-dependent changes in cell motility. The gene encoding human Fe65 maps to chromosome 11p15.

REFERENCES

- Duilio, A., et al. 1991. A rat brain mRNA encoding a transcriptional activator homologous to the DNA binding domain of retroviral integrases. Nucleic Acids Res 19: 5269-5274.
- 2. Bressler, S.L., et al. 1996. cDNA cloning and chromosome mapping of the human Fe65 gene: interaction of the conserved cytoplasmic domains of the human β -Amyloid precursor protein and its homologues with the mouse Fe65 protein. Hum. Mol. Genet. 5: 1589-1598.
- Borg, J.P., et al. 1996. The phosphotyrosine interaction domains of X11 and Fe65 bind to distinct sites on the YENPTY motif of amyloid precursor protein. Mol. Cell. Biol. 16: 6229-6241.
- Guenette, S.Y., et al. 1999. hFe65L influences amyloid precursor protein maturation and secretion. J. Neurochem. 73: 985-993.
- Sabo, S.L., et al. 1999. Regulation of β-amyloid secretion by FE65, an amyloid protein precursor-binding protein. J. Biol. Chem. 274: 7952-7957.
- Ando, K., et al. 2001. Phosphorylation-dependent regulation of the interaction of amyloid precursor protein with Fe65 affects the production of β-Amyloid. J. Biol. Chem. 276: 40353-40361.

CHROMOSOMAL LOCATION

Genetic locus: APBB1 (human) mapping to 11p15.4; Apbb1 (mouse) mapping to 7 E3.

SOURCE

Fe65 (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Fe65 of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-19750 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Fe65 (N-20) is recommended for detection of Fe65 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Fe65 (N-20) is also recommended for detection of Fe65 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Fe65 siRNA (h): sc-41954, Fe65 siRNA (m): sc-41955, Fe65 shRNA Plasmid (h): sc-41954-SH, Fe65 shRNA Plasmid (m): sc-41955-SH, Fe65 shRNA (h) Lentiviral Particles: sc-41954-V and Fe65 shRNA (m) Lentiviral Particles: sc-41955-V.

Molecular Weight of Fe65: 85-90 kDa.

Positive Controls: Fe65 (h) 293T Lysate: sc-175237, NIH/3T3 whole cell lysate: sc-2210 or HeLa whole cell lysate: sc-2200.

DATA



Fe65 (N-20): sc-19750. Western blot analysis of Fe65 expression in non-transfected: sc-117752 (**A**) and human Fe65 transfected: sc-175237 (**B**) 293T whole cell lysates.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfation
GuaranteedTry Fe65 (F-6): sc-398389 or Fe65 (D-11): sc-374641,
our highly recommended monoclonal alternatives to
Fe65 (N-20).