

GGT1 (T-20): sc-19798

BACKGROUND

GGT (γ -glutamyltranspeptidase) acts as a glutathionase and catalyzes the transfer of the glutamyl moiety of glutathione to a variety of amino acids and dipeptide acceptors. This enzyme is located on the outer surface of the cell membrane and is widely distributed in mammalian tissues involved in absorption and secretion. In humans, hepatic GGT activity is elevated in some liver diseases. GGT1 is released into the bloodstream after liver damage, and an elevated level of the enzyme may be a useful early sign of hepatocellular carcinoma. GGT5 converts leukotriene C4 to leukotriene D4; it does not, however, convert synthetic substrates that are commonly used to assay GGT. In human serum and in human tissues, there is a marked heterogeneity in GGT, but this heterogeneity can be attributed to different glycosylation of the same peptide rather than to the products of different genes.

REFERENCES

1. Bulle, F., Mattei, M.G., Siegrist, S., Pawlak, A., Passage, E., Chobert, M.N., Laperche, Y. and Guellaen, G. 1987. Assignment of the human γ -glutamyl transferase gene to the long arm of chromosome 22. *Hum. Genet.* 76: 283-286.
2. Heisterkamp, N., Rajpert-De Meyts, E., Uribe, L., Forman, H.J. and Groffen, J. 1991. Identification of a human γ -glutamyl cleaving enzyme related to, but distinct from, γ -glutamyl transpeptidase. *Proc. Natl. Acad. Sci. USA* 88: 6303-6307.
3. Visvikis, A., Thioudellet, C., Oster, T., Fournel-Gigleux, S., Wellman, M. and Siest, G. 1991. High-level expression of enzymatically active mature human γ -glutamyltransferase in transgenic V79 Chinese hamster cells. *Proc. Natl. Acad. Sci. USA* 88: 7361-7365.
4. LocusLink Report (LocusID: 137181). <http://www.ncbi.nlm.nih.gov/LocusLink>

SOURCE

GGT1 (T-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of GGT1 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-19798 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

GGT1 (T-20) is recommended for detection of GGT1 heavy chain of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GGT1 siRNA (m): sc-35474, GGT1 shRNA Plasmid (m): sc-35474-SH and GGT1 shRNA (m) Lentiviral Particles: sc-35474-V.


Molecular Weight of GGT1: 64 kDa.

Positive Controls: mouse kidney extract: sc-2255 or rat kidney extract: sc-2394.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



GGT1 (T-20): sc-19798. Western blot analysis of GGT1 expression in mouse kidney tissue extract.



GGT1 (T-20): sc-19798. Western blot analysis of GGT1 expression in non-transfected (A) and mouse GGT1 transfected: sc-120476 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **GGT1 (F-7): sc-374495** or **GGT1 (3E6): sc-100746**, our highly recommended monoclonal alternatives to GGT1 (T-20).