



PEPT2 (C-20): sc-19920

BACKGROUND

In mammalian small intestine, the proton-coupled peptide transporter (PEPT) is responsible for the absorption of small peptides arising from digestion of dietary proteins. PEPT1, a hydrogen ion/peptide co-transporter, transports dipeptides and tripeptides, but not free amino acids or peptides with more than three amino acid residues. The mechanism for this transport requires proton binding and the presence of an inside-negative membrane potential. PEPT1 is 708 amino acid protein that contains 12 putative membrane-spanning domains. PEPT1 seems to play important roles in nutritional and pharmacological therapies. The mammalian kidney expresses a proton-coupled peptide transporter, PEPT2, that is responsible for the absorption of small peptides, as well as beta-lactam antibiotics and other peptide-like drugs, from the tubular filtrate.

REFERENCES

- Liang, R., Fei, Y.J., Prasad, P.D., Ramamoorthy, S., Han, H., Yang-Feng, T.L., Hediger, M.A., Ganapathy, V. and Leibach, F.H. 1995. Human intestinal H⁺/peptide cotransporter. Cloning, functional expression, and chromosomal localization. *J. Biol. Chem.* 270: 6456-6463.
- Liu, W., Liang, R., Ramamoorthy, S., Fei, Y.J., Ganapathy, M.E., Hediger, M.A., Ganapathy, V. and Leibach, F.H. 1995. Molecular cloning of PEPT2, a new member of the H⁺/peptide cotransporter family, from human kidney. *Biochim. Biophys. Acta* 1235: 461-466.
- Adibi, S.A. 1997. The oligopeptide transporter (PEPT1) in human intestine: biology and function. *Gastroenterology* 113: 332-40.
- Ocheltree, S.M., Shen, H., Hu, Y., Xiang, J., Keep, R.F. and Smith, D.E. 2004. Role of PEPT2 in the choroid plexus uptake of glycylsarcosine and 5-aminolevulinic acid: studies in wild-type and null mice. *Pharm. Res.* 21:1680-1685.
- Meredith, D. and Price, R.A. 2006. Molecular modeling of PEPT1--towards a structure. *J. Membr. Biol.* 213: 79-88.
- Terada, T. and Inui, K. 2007. Gene expression and regulation of drug transporters in the intestine and kidney. *Biochem. Pharmacol.* 73: 440-449.

CHROMOSOMAL LOCATION

Genetic locus: SLC15A2 (human) mapping to 3q13.33; Slc15a2 (mouse) mapping to 16 B3.

SOURCE

PEPT2 (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of PEPT2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-19920 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

PEPT2 (C-20) is recommended for detection of PEPT2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PEPT2 siRNA (h): sc-42590.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.